

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 565 567 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: **18.01.95** (51) Int. Cl.⁸: **C07K 7/56, C07K 5/06, A61K 35/66**

(21) Application number: **92901892.7**

(22) Date of filing: **02.01.92**

(96) International application number:
PCT/EP92/00002

(97) International publication number:
WO 92/12172 (23.07.92 92/19)

Consolidated with 92100025.3/0494078
(European application No./publication No.) by
decision dated 21.02.94.

(94) **AMIDES OF ANTIBIOTIC GE 2270 FACTORS.**

(30) Priority: **03.01.91 EP 91100123**

(43) Date of publication of application:
20.10.93 Bulletin 93/42

(45) Publication of the grant of the patent:
18.01.95 Bulletin 95/03

(94) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU MC
NL PT SE**

(73) Proprietor: **GRUPPO LEPETIT S.p.A.**
Via Roberto Lepetit, 8
I-20020 Lainate (MI) (IT)

(72) Inventor: **TAVECCHIA, Paolo**
Via Buon Gesù, 1
I-20017 Rho (IT)
Inventor: **LOCIURO, Sergio**
Via Troilo, 4

I-20136 Milano (IT)
Inventor: **CIABATTI, Romeo**
Via Brodolini, 15/A
I-20026 Novate Milanese (IT)
Inventor: **SELVA, Enrico**
Via Di Vittorio, 15
I-27027 Gropello Cairoli (IT)

(74) Representative: **Macchetta, Francesco et al**
GRUPPO LEPETIT S.p.A.
Patent and Trademark Department
Via Roberto Lepetit, 34
I-21040 Gerenzano (Varese) (IT)

EP 0 565 567 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

(C₁-C₄)alkyl,
 amino(C₂-C₄)alkyl,
 (C₁-C₄)alkylamino-(C₁-C₄)alkyl, or
 di-(C₁-C₄)alkylamino-(C₁-C₄)alkyl;
 5 R₃ represents:
 hydrogen,
 a linear or branched (C₁-C₁₄)alkyl group bearing from 1 to 3 substituents selected from:
 carboxy, sulfo, phosphono, amino which may be optionally protected with a lower
 alkoxy carbonyl or a benzyloxycarbonyl group, (C₁-C₄)alkylamino wherein the alkyl
 10 moiety may be optionally substituted with a carboxy group,
 di-(C₁-C₄)alkylamino, hydroxy, halo,
 (C₁-C₄)alkoxy wherein the alkyl moiety may be optionally substituted with a carboxy
 group,
 (C₁-C₄)alkoxy carbonyl, mercapto,
 15 (C₁-C₄)alkylthio wherein the alkyl moiety may be optionally substituted with a carboxy
 group, phenyl which may be optionally substituted with 1 to 3 substituents selected
 from carboxy, sulfo, hydroxy, halo and mercapto, carbamyl,
 (C₁-C₆)alkylcarbamyl wherein the alkyl moiety may be optionally substituted with 1 or 2
 substituents selected from carboxy, amino,
 20 (C₁-C₄)alkylamino and di-(C₁-C₄)alkylamino,
 di-(C₁-C₄)alkylcarbamyl wherein the alkyl moieties together with the adjacent nitrogen
 atom may also represent a saturated 5-7 membered heterocyclic ring which may
 optionally be substituted with a carboxy or a carbamyl group on one of the ring carbons
 and may optionally contain a further heterogroup selected from O, S and N, ben-
 25 zoylamino wherein the phenyl group may be substituted from 1 to 3 hydroxy group, a
 nitrogen containing 5-6 membered heterocyclic ring which may be unsaturated, partially
 saturated or wholly saturated and may contain 1 to 3 further heteroatoms selected from
 N, S and O wherein one of the carbons of the ring may optionally bear a group
 carboxy, sulfo,
 30 carboxy(C₁-C₄)alkyl or sulfo(C₁-C₄)alkyl and the ring nitrogen atom may optionally be
 substituted by (C₁-C₄)alkyl,
 carboxy(C₁-C₄)alkyl, sulfo(C₁-C₄)alkyl, or benzyl;
 (C₃-C₆)alkenyl, optionally substituted by carboxy or sulfo;
 1-deoxy-1-glucityl;
 35 2-deoxy-2-glucosyl;
 a fully saturated 5 to 7 membered nitrogen containing heterocyclic ring wherein the
 nitrogen atom may be optionally substituted by (C₁-C₄)alkyl or benzyl and one or two
 carbons of the ring skeleton may bear a substituent selected from (C₁-C₄)alkyl, carboxy
 and sulfo;
 40 or R₂ and R₃ taken together with the adjacent nitrogen atom represent a fully saturated 5-7 mem-
 bered heterocyclic ring which may optionally contain a further heteroatom selected
 from O, S and N, and may optionally bear one or two substituents on the ring carbons
 selected from (C₁-C₄)alkyl, benzyl, carboxy, sulfo, carboxy(C₁-C₄)alkyl, and sulfo(C₁-
 C₄)alkyl;
 45 R₄ represents:
 hydrogen,
 methyl, or
 hydroxymethyl;


with the proviso that when R₄ is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and

50 R₁ is methyl;

and the pharmaceutically acceptable addition salts thereof.

This invention includes also a process for preparing the compounds of this invention from the
 corresponding starting compounds of formula (II)

55



wherein W is a carboxylic function or an activated ester thereof, and R, R₁ and R₄ being as defined above.


Antibiotic GE 2270 is prepared by culturing a sample of *Planobispora rosea* ATCC 53773 or a producing variant or mutant thereof and isolating the desired antibiotic substance from the mycelium and/or the fermentation broth. *Planobispora rosea* ATCC 53773 was isolated from a soil sample and deposited on June 14, 1988 with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, MD 20852 Maryland, U.S.A., under the provisions of the Budapest Treaty.

The strain has been accorded accession number ATCC 53773.

Antibiotic GE 2270 factor A is the main component of the antibiotic GE 2270 complex.

Antibiotic GE 2270 factor A and *Planobispora rosea* ATCC 53773 are described in European Patent Application Publication No. 359062.

Recent studies showed that antibiotic GE 2270 factor A can be represented by the following general formula III



When antibiotic GE 2270 factor A is treated under selective hydrolysis conditions some derivatives named antibiotic GE 2270 factors A₁, A₂ and A₃ are obtained. Said factors A₁, A₂ and A₃ and the hydrolysis process for preparing them are disclosed in the European Patent Application Publication No. 406745 and U.S. Patent Application No. 547,647.


Generally, the above mentioned hydrolytic conditions involve the use of mixtures of buffered or unbuffered aqueous acid media and polar organic solvents. The reaction temperature varies depending on factors such as the strength and the concentration of the acid employed, and is generally comprised between -10 °C and 90 °C. Also the reaction time varies considerably depending on parameters such as the temperature, the acid strength and its concentration; generally, it may vary from a few minutes to several hours.

In general, when milder hydrolysis conditions are employed, e.g. shorter reaction time and lower temperature or lower acid strength or concentration, antibiotic GE 2270 factor A₁ is normally obtained, while stronger hydrolysis conditions yield antibiotic GE 2270 factor A₂. To obtain antibiotic GB 2270 factor A₃, still more drastic hydrolysis conditions are necessary.

While antibiotic GE 2270 factors A₂ and A₃ can be directly utilized as the starting materials for the production of the compounds of this invention, antibiotic GE 2270 factor A₁ is not suitable as the starting material for direct production of the compounds of this invention; however, it can be utilized as a precursor of the said starting materials as it will be explained further.

Antibiotic GE 2270 factor A₂ and factor A₃ are characterized by having an ester and a carboxy function respectively in the upper part of the molecule. In particular, it has been found that antibiotic GE 2270 factor A₂ and factor A₃ can be represented by the above defined formula II wherein:

W represents COOH (antibiotic GE 2270 factor A₃) or the ester moiety (antibiotic GE 2270 factor A₂)



R is methoxymethyl,

R₁ is methyl and

R₄ is methyl.

Both antibiotic GE 2270 factor A₂ and factor A₃ (and a mixture thereof) can be used as suitable starting materials for the production of the compounds of the invention, even if factor A₃ is the preferred one. Factor A₂ may be employed directly as an activated ester or may be converted to factor A₃ by drastic acid hydrolysis conditions, as mentioned above, or by basic hydrolysis with diluted alkali (as described in European Patent Application Publication No. 406745 and U.S. Patent Application No. 547,647).

It was recently found (European Patent Application Publication No. 451486 and U.S. Patent Application No. 665,612) that other minor components can be isolated from the cultures of *Planobispora rosea* ATCC 53773 or an antibiotic GE 2270 producing variant or mutant thereof. In particular, they are found in the mycelium and also in the fermentation broths of the cultured microorganism.

A preferred procedure for recovering said minor components of antibiotic GE 2270 from the mycelium includes extracting the filtered or centrifugated mycelium with a water-miscible organic solvent, concentrating the extracts and recovering the crude antibiotic substance by precipitation, optionally with the addition of a precipitating agent, by extraction of the aqueous residue with a water-immiscible organic solvent or by adsorption chromatography followed by elution of the desired product from the absorption matrix.

It was recently found (European Patent Application No. 91114667.8) that a further minor component (factor C_{2a}) can be isolated from the same culture of *Planobispora rosea* ATCC 53773 described above.

The physico-chemical characteristics of antibiotic GE 2270 C_{2a} are the following:

A) The ultraviolet absorption spectrum recorded with a Perkin Elmer Model 320 spectrometer exhibits the following absorption maxima:

Solvent	UV max (nm)
0.1 M HCl	245-250 (shoulder) 300-315
0.1 M KOH	245-250 (shoulder) 300-315
Phosphate buffer pH 7.38	245-250 (shoulder) 300-315
Methanol	245-250 (shoulder) 300-315

B) The ¹H-NMR spectrum of antibiotic GE 2270 factor C_{2a} was recorded at 250 MHz with a Bruker spectrometer. The spectrum of the antibiotic in DMSO-d₆ (hexadeuterodimethylsulfoxide) using TMS as the internal standard (δ 0.00 ppm) exhibits the following groups of signals [δ, ppm, multiplicity] (s = singlet, d = doublet, t = triplet, m = multiplet, Py = pyridine, Tz = thiazole):

9.03, d, (NH); 8.70, d, (2NH's); 8.60, s, 8.54, s, 8.29, s, and 7.38, s, (Tz CH's); 8.48, m, (glycine NH); 8.43, d, and 8.27, d, (Py CH's); 7.35-7.20, m, (aromatic CH's and primary amide NH); 6.98, s (primary amide NH); 6.04, d, (OH); 5.80, t (OH); 5.35-5.15, m, (αCH's); 5.04, m, (phenylserine βCH); 4.98, s [CH₂-(OCH₃)]; 4.87, d, [CH₂(OH)]; 4.81, m and 4.56, m, (oxazoline CH₂); 4.35-3.75, m, (CH₂ of glycine and prolineamide CH's); 3.39, s, (OCH₃); 2.71, m, and 1.30, m, (CH₂ of asparagine); 2.48, d, (NCH₃ of N-methylasparagine); 2.22-1.80, m, (isopropyl CH and prolineamide CH's); 0.88 and 0.84, d, (valine CH₃'s)

C) Antibiotic GE 2270 factor C_{2a} shows retention time (R_t) of 12.6 min and retention time relative to antibiotic GE 2270 factor A (R_t 16.6 min) of 0.76 when analyzed with the following reverse phase HPLC

system:

Column: Bakerbond® C8 (5 µm) 4.6x250 mm (Bakerbond® is a trade name for reverse phase octylsilyl silica gel HPLC columns supplied by J.T. Baker Research Product, Phillipsburg, New Jersey 08865 USA)

Flow rate: 1.8 ml/min

5 Phase A: CH₃CN:tetrahydrofuran:40 mM HCOONH₄ 40:40:20 (v/v/v)

Phase B: CH₃CN:tetrahydrofuran:40 mM HCOONH₄ 10:10:80 (v/v/v)


Elution: linear gradient from 20% to 30% of Phase A in 20 min

Detection: UV 254 nm

10 D) The main FAB-MS peak of antibiotic GE 2270 factor C_{2a} is 1306 daltons. This corresponds most likely to the lowest isotope of the protonated molecular ion. The analysis was performed on a Kratos MS-50 double focusing mass spectrometer, using 8 kV accelerating voltage and a saddle field atom gun with Xe gas (2x10⁻⁵ torr pressure indicated on the source ion gauge) at 6 kV voltage and 1 mA current. The antibiotic for the FAB-MS analysis was mixed with a thioglycerol matrix containing 0.1 M acetic acid.

15 Some of said minor components of antibiotic GE 2270 (i.e. factors B₁, B₂, C₁, C₂, C_{2a}, D₁, D₂ and E) may be represented by the general formula II mentioned above wherein

W represents the moiety:



30 R represents respectively hydrogen for GE 2270 factors C₁ and D₁, methyl for factor B₂, hydroxymethyl for factors D₂ and E and methoxymethyl for factors B₁, C₂ and C_{2a};

R₁ represents hydrogen for GE 2270 factors B₁, D₁ and E and methyl for GE 2270 factors B₂, C₁, C₂, C_{2a} and D₂; and

35 R₄ represents hydrogen for GE 2270 factor C₂, methyl for GE 2270 factors B₁, B₂, C₁, D₁, D₂ and E and hydroxymethyl for factor C_{2a}.

When antibiotic GE 2270 factors D₁, D₂ and E or a mixture thereof are treated by the same hydrolytic process outlined above (and described in European Patent Application Publication No. 406745 and U.S. Patent Application No. 547,647) for preparing antibiotic GE 2270 factors A₂ and A₃ from antibiotic GE 2270 factor A, the common moiety W cited above is hydrolyzed to a carboxy moiety leaving the substituents R, R₁ and R₄ unaltered.

40 Therefore, the derivatives of formula II wherein W is a carboxy or an activated ester function, R is hydrogen, hydroxymethyl or methoxymethyl, R₁ is hydrogen or methyl and R₄ is hydrogen, methyl or hydroxymethyl, with the proviso that when R₄ is hydrogen or hydroxymethyl then R is methoxymethyl and R₁ is methyl, can be used as starting materials of the present invention. It has to be clear that as with other microorganisms, the characteristics of the GE 2270 producing strains are subject to variation. For example, artificial variants and mutants of the strain can be obtained by treatment with various known mutagens, such as U.V. rays, X-rays, high frequency waves, radioactive rays, and chemicals such as nitrous acid; N-methyl-N'-nitro-N-nitroso-guanidine, and many others. All natural and artificial variants and mutants which belong to a species of the genus *Planobispora* and produce antibiotic GE 2270 are deemed equivalent to strain

50 *Planobispora rosea* ATCC 53773 for the purposes of this invention.

As used herein, the term "alkyl", either alone or in combination with other substituents, includes both straight and branched hydrocarbon groups; more particularly, "(C₁-C₁₄)alkyl" represents a straight or branched aliphatic hydrocarbon chain of 1 to 14 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 1-hexyl, 2-hexyl, 3-hexyl, 3,3-dimethyl-1-butyl, 4-methyl-1-pentyl and 3-methyl-1-pentyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tridecyl and tetradecyl; likewise, "(C₁-C₄)alkyl" represents a straight or branched hydrocarbon chain of 1 to 4 carbon atoms such as those alkyl of 1 to 4 carbons exemplified above.

As described above the "(C₁-C₁₄)alkyl" moiety may bear 1 to 3 substituents.


The term "halo" represents a halogen atom selected from fluoro, chloro, bromo and iodo.

As used herein, the term "(C₃-C₆)alkenyl" means an alkylene radical having three to six carbon atoms and a double bond; it comprises propenyl, 3-butenyl, 2-butenyl, 2-methylpropenyl, 2-pentenyl, 3-hexenyl and so on, which may be optionally substituted with a carboxy or a sulfo group.

The expression "a nitrogen containing 5-6 membered heterocyclic ring which may contain 1 to 3 further heteroatoms selected from N, S and O" according to the present invention includes unsaturated, partially saturated and wholly saturated ring systems such as pyridine, pyrimidine, pyrazine, pyrrolidine, piperidine, piperazine, oxazole, oxazoline, oxazolidine, pyrazoline, pyrazolidine, thiazolidine, morpholine, thiomorpholine, pyrrole, pyrroline, imidazole, imidazolidine, thiadiazole, oxadiazole and tetrazole.

In said "nitrogen containing 5-6 membered heterocyclic ring" 1 to 3 ring carbons may optionally bear a group carboxy, sulfo, carboxy(C₁-C₄)alkyl and sulfo(C₁-C₄)alkyl and the ring nitrogen atom may optionally be substituted by (C₁-C₄)alkyl, carboxy(C₁-C₄)alkyl, sulfo(C₁-C₄)alkyl, or benzyl.

The expression "fully saturated 5-7 membered nitrogen containing heterocyclic ring wherein the nitrogen atom may be optionally substituted by (C₁-C₄)alkyl or benzyl" identifies a fully saturated heterocycle ring of 5-7 members containing a nitrogen atom which can be optionally substituted by (C₁-C₄)alkyl or benzyl wherein the carbon skeleton may optionally bear one or two substituents selected from (C₁-C₄)alkyl, carboxy and sulfo. Said heterocyclic rings are connected with the nitrogen atom of the group:




through a bond between the same nitrogen moiety and a carbon atom of the heterocyclic rest. Examples of said radicals are: 1-methyl-4-pyrrolidinyl, 3-piperidinyl, 1-ethyl-4-piperidinyl, 1-benzyl-2,6-dimethyl-4-piperidinyl, and 4-carboxy-1-methyl-2-piperidinyl.

When R₂ and R₃ taken together with the adjacent nitrogen atom represent "a fully saturated 5-7 membered heterocyclic ring which may optionally contain a further heteroatom selected from O, S and N" this expression includes, for instance, the following heterocyclic groups: pyrrolidino, morpholino, piperidino, piperazino, thiomorpholino, pyrazolidino, 1,3-oxazolidino, 1,3-thiazolidino and hexahydroazepino. When the further heteroatom is N it may optionally bear a substituent selected from (C₁-C₄)alkyl, benzyl, carboxy, carboxy(C₁-C₄)alkyl, sulfo and sulfo(C₁-C₄)alkyl.

The term "1-deoxy-1-glucityl" identifies a compound of formula (I) wherein Y is a radical deriving from glucamine, i.e. 1-amino-1-deoxy-glucitol. The term "2-deoxy-2-glucosyl" identifies a compound of formula (I) wherein Y is a radical deriving from glucosamine, i.e. 2-amino-2-deoxyglucose.

A preferred group of compounds of the invention is represented by those compounds of formula I wherein R represents methoxymethyl, R₁ and R₄ represent a methyl group and the other substituents are as defined above.

A further preferred group of compounds of the invention are those compounds of formula I wherein R represents methoxymethyl, R₁ and R₄ represent a methyl group, and Y represents a group of formula



wherein R₂ is hydrogen and R₃ is defined as above.


A further preferred group of compounds of the invention is represented by those compounds of formula I wherein R is methoxymethyl, R₁ and R₄ represent a methyl group and Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.

A further preferred group of compounds comprises those compounds of formula I wherein R is methoxymethyl, R_1 and R_4 are methyl, Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH, SO_3H and PO_3H_2 .

5 The most preferred compound is represented by the formula I wherein R is methoxymethyl, R_1 and R_4 are methyl and Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is $CH_2CH_2CH_2CH_2CH_2-COOH$.

A further preferred group of compounds of the invention are those compounds of formula I wherein R represents hydrogen, hydroxymethyl or methoxymethyl, R_1 represents hydrogen or a methyl group, and Y represents a group of formula

10



15

wherein R_2 is hydrogen and R_3 and R_4 are defined as above.


20 A further preferred group of compounds of the invention is represented by those compounds of formula I wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 represents hydrogen or a methyl group, R_4 is hydrogen, methyl or hydroxymethyl with the proviso that when R_4 is hydrogen or hydroxymethyl then R is methoxymethyl and R_1 is methyl, and Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.

25 A further preferred group of compounds comprises those compounds of formula I wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 is hydrogen or methyl, R_4 is hydrogen, methyl or hydroxymethyl with the proviso that when R_4 is hydrogen or hydroxymethyl then R is methoxymethyl and R_1 is methyl, Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH, SO_3H and PO_3H_2 .

30 The last preferred group of compounds is represented by those compounds of the formula I wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 is hydrogen or methyl, R_4 is as defined above and Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is $CH_2CH_2CH_2CH_2CH_2-COOH$.

35 Representative examples of the compounds of the invention, include those compounds of formula I wherein R, R_1 , R_4 and Y are as defined above and

40



45

represents

-NH₂

-NHC₄H₉

50 -NH(CH₂)₄-PO₃H₂


-NHCH₂COOH

-NH-CH₂CONH₂


-NH-CH₂-CON(C₂H₅)₂

55


5




10




15




20



25








30



35

wherein n is 2, 3 or 4

40




45

wherein n is 2, 3, 4, 5, 6, 7 or 8


50

55

5



10



15

20

25

30








35






40

45


50

55






5



10

15


20



25

30

35




40

45

50


55

5




10

15




20

25




30



35





40









45

50

55





wherein n is 2, 3, 4 or 5 and m is 1, 2, 3 or 4

$\text{-NH-(CH}_2\text{)}_n\text{-CH = CH-(CH}_2\text{)}_m\text{-COOH}$

$\text{-NH-(CH}_2\text{)}_n\text{-CH = CH-(CH}_2\text{)}_m\text{-SO}_3\text{H}$

wherein n is 1, 2 or 3 and m is 0, 1 or 2

5 The compounds of the invention can form salts according to conventional procedures.

In particular, those compounds of formula I wherein the group $\text{-NR}_2\text{R}_3$ contains further amine functions form acid addition salts.

In addition, those compounds of the invention which contain acid functions in the $\text{-NR}_2\text{R}_3$ moiety may also form base addition salts.

10 In general, those compounds of the invention which contain acid and basic functions can form internal salts. For the scope of the present invention the "internal salts" are encompassed by the definition of the "non-salt" form.

Preferred addition salts of the compounds of this invention are the pharmaceutically acceptable acid and/or base addition salts.

15 With the term "pharmaceutically acceptable acid and/or base addition salts" are intended those salts with acids and/or bases which from biological, manufacturing and formulation standpoint are compatible with the pharmaceutical practice as well as with the use in the animal growth promotion.

Representative and suitable acid addition salts of the compounds of formula I include those salts formed by standard reaction with both organic and inorganic acids such as, for example, hydrochloric, 20 hydrobromic, sulfuric, phosphoric, acetic, trifluoroacetic, trichloroacetic, succinic, citric, ascorbic, lactic, maleic, fumaric, palmitic, cholic, pantoic, mucic, glutamic, camphoric, glutaric, glycolic, phthalic, tartaric, lauric, stearic, salicylic, methanesulfonic, dodecylsulfonic (estolic acid), benzenesulfonic, sorbic, picric, benzoic, cinnamic and the like acids.

Representative examples of these bases are:

25 alkali metal or alkaline-earth metal hydroxides such as sodium, potassium, and calcium hydroxide; ammonia and organic aliphatic, alicyclic or aromatic amines such as methylamine, dimethylamine, trimethylamine, 2-amino-2-hydroxymethyl-1,3-propanediol (TRIS), picoline and basic aminoacids such as lysine, ornithine, arginine and histidine.

30 The transformation of the free amino or non-salt compounds of the invention into the corresponding addition salts, and the reverse, i.e. the transformation of an addition salt of a compound of the invention into the non-salt or free amino form, are within the ordinary technical skill and are encompassed by the present invention.

For instance, a compound of formula I can be transformed into the corresponding acid or base addition-salt by dissolving the non-salt form in an aqueous solvent and adding a slight molar excess of the selected 35 acid or base. The resulting solution or suspension is then lyophilized to recover the desired salt. Instead of lyophilizing, in some instances, it is possible to recover the final salt by extraction with an organic solvent, concentration to a small volume of the separated organic phase and precipitation by adding a non-solvent.

In case the final salt is insoluble in an organic solvent where the non-salt form is soluble it is recovered by filtration from the organic solution of the non-salt form after addition of the stoichiometric amount or a 40 slight molar excess of the selected acid or base.

The non-salt form can be prepared from a corresponding acid or base salt dissolved in an aqueous solvent which is then neutralized to free the non-salt form. This is then recovered for instance by extraction with an organic solvent or is transformed into another base or acid addition salt by adding the selected acid or base and working up as above.

45 When following the neutralization desalting is necessary, a common desalting procedure may be employed.

For example, column chromatography on controlled pore polydextrane resins (such as Sephadex LH 20) or silanized silica gel may be conveniently used. After eluting the undesired salts with an aqueous solution, the desired product is eluted by means of linear gradient or step-gradient of a mixture of water and 50 a polar or apolar organic solvent, such as acetonitrile/water from 50:50 to about 100% acetonitrile.

As is known in the art, the salt formation either with pharmaceutically acceptable acids (bases) or non-pharmaceutically acceptable acids (bases) may be used as a convenient purification technique. After formation and isolation, salt form of a compound of formula I can be transformed into the corresponding non-salt or into a pharmaceutically acceptable salt.

55 In some instances the acid addition salt of a compound of formula I is more soluble in water and hydrophilic solvents and has an increased chemical stability.


However, in view of the similarity of the properties of the compounds of formula I and their salts, what is said in the present application when dealing with the biological activities of the compounds of formula I

applies also to their pharmaceutically acceptable salts, and viceversa.

In view of their properties, the compounds of the invention can be used as active ingredients in the preparation of medicaments for human or animal treatment.

In particular, the amide derivatives of the antibiotic GE 2270 compounds of formula I are antimicrobial agents mainly active against gram positive bacteria and gram positive as well as gram negative anaerobes.

A general procedure for preparing a compound of this invention is represented by the reaction (amidation) of a suitable antibiotic GE 2270 compound having formula II



wherein

- 35 W represents a carboxy or an activated ester function;
 R represents hydrogen, hydroxymethyl or methoxymethyl;
 R₁ represents hydrogen or methyl;
 R₄ represents hydrogen, methyl or hydroxymethyl,

with the proviso that, when R₄ represents hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and R₁ is methyl; with a selected amino of formula HNR₂R₃ wherein R₂ and R₃ have the same meanings as above in an inert organic solvent and, when W is carboxy, in the presence of a condensing agent.

In carrying out the amidation for preparing the compounds of this invention, sometimes, it is convenient to protect the functions of the reactants which are not involved in the amidation reaction but could result sensitive to the reaction conditions or negatively affect the reaction course, for instance, yielding undesired side-product.

Furthermore, when the amino acid contains further reactive functions such as amino, carboxy or mercapto groups which may interfere with the course of the amidation, these are protected by means of methods known per se in the art such as those described in reference books like E. Gross and J. Meienhofer "The Peptides", Vol. 3, Academic Press, New York, 1981 and M. Bodanszky and A. Bodanszky "The Practice of Peptide Synthesis", Springer-Verlag, Berlin, Heidelberg, 1984. These protecting groups must be stable at the conditions the amidation reaction takes place and must be easily removable at the end of the reaction without affecting either the newly formed amide bond or any other part of the molecule.

Representative examples of N-protecting groups which may be advantageously used in the process of the invention for protecting an amino function are carbamate forming reagents characterized by the following oxycarbonyl groups: 1,1-dimethylpropynyloxycarbonyl, t-butyloxycarbonyl, vinyloxycarbonyl, arylloxycarbonyl, cinnamyloxycarbonyl, benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 3,4-dimethoxy-6-nitrobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 5-benzisoxazolylmethyloxycarbonyl, 9-anthranyl-

methyloxycarbonyl, diphenylmethyloxycarbonyl, isonicotinylloxycarbonyl, S-benzyloxycarbonyl, and the like.

A suitable protection for reactive carboxylic acid function is, for instance, by forming an ester function.

The man skilled in the art is capable, also on the basis of the present disclosure, of deciding which functions of the amino HNR_2R_3 need to be protected, how they must be protected and the proper deprotection reaction which is necessary to free the final compound.

As it is appreciated by the skilled technician, the ultimate choice of the specific protecting group depends on the characteristics of the particular amide derivative which is desired. In fact, this amide function of the final compound should be stable at the condition of removal of the protecting group(s).

Since the conditions of removal of the different protecting groups are known, the skilled technician is capable of selecting the proper protecting group.

Inert organic solvents useful for the condensation reaction are those solvents which do not unfavorably interfere with the reaction course and are capable of at least partially solubilizing the antibiotic starting material.

Examples of said inert solvents are organic amides, ethers of glycols and polyols, phosphoramides, sulfoxides. Preferred examples of inert solvents are: dimethylformamide, dimethoxyethane, hexamethylphosphoramide, dimethylsulfoxide, dioxane, and mixtures thereof.

Sometimes, water is compatible with the reaction conditions.

The condensing agent in the process of the invention when W is carboxy is one suitable for forming amide bonds in organic compounds and in particular in peptide synthesis.

Representative and preferred examples of condensing agents are $(\text{C}_1\text{-C}_4)$ alkyl, phenyl or heterocyclic phosphorazidates such as, diphenylphosphorazidate (DPPA), diethylphosphorazidate, di(4-nitrophenyl)-phosphorazidate, dimorpholylphosphorazidate and diphenylphosphorochloridate or benzotriazol-1-yl-oxy-trispyrrolidinophosphoniumhexafluorophosphate (PyBOP). The preferred condensing agent is diphenylphosphorazidate (DPPA).

In the process of the invention, the amine reactant HNR_2R_3 is normally used in a slight molar excess.

In general, a 1- to 2-fold molar excess is used while a 1.2- to 1.5-fold molar excess is preferred.

For the amidation to proceed, it is necessary that the amine HNR_2R_3 be capable of forming a salt with the carboxy function of the antibiotic starting material. In case the amine HNR_2R_3 is not strong enough to form such a salt in the selected reaction medium, it is necessary to add a salt-forming base to the reaction mixture at least in an equimolecular amount with the antibiotic starting material.

Examples of said salt-forming bases are tertiary organic aliphatic or alicyclic amines such as trimethylamine, triethylamine, N-methyl pyrrolidine or heterocyclic bases such as picoline, and the like.

The condensing agent is generally employed in a slight molar excess such as from 1.1 to 1.5 and preferably is 1.2 times the antibiotic GE 2270 starting compound.

In addition, the amine reactant HNR_2R_3 may also conveniently be introduced in the reaction medium as a corresponding acid addition salt, e.g. the hydrochloride. In this case, at least a double molar proportion and preferably a 2 to 3 fold molar excess of a strong base capable of freeing the HNR_2R_3 amine from its salts, is used. Also in this case, the suitable base is a tertiary organic aliphatic or alicyclic amine like those exemplified above. In fact, at least in some instances, the use of salt of the amine HNR_2R_3 , which is then freed *in situ* with the above mentioned bases, is greatly preferred especially when the salt is more stable than the corresponding free amine.

The reaction temperature will vary considerably depending on the specific starting materials and reaction conditions. In general, it is preferred to conduct the reaction at temperatures between 0-20 °C.

Also the reaction time vary considerably depending on the other reaction parameters. In general the condensation reaction is completed in about 5-24 h.

In any case, the reaction course is monitored by TLC or preferably by HPLC according to methods known in the art.

On the basis of the results of these assays a man skilled in the art will be able to evaluate the reaction course and decide when to stop the reaction and start working up the reaction mass according to known *per se* techniques which include, for instance, extraction with solvents, precipitation by addition of non-solvents, etc., in conjunction with further separations and purifications by column chromatography.

As already said, when protection of the HNR_2R_3 reactant is necessary, the protected final compound is then de-protected according to procedures which are known *per se* and mainly depend on the protecting group involved.

When an activated ester of GE 2270 is used as the starting material, said ester is one wherein the esterified alcohol is providing a leaving group which can be readily displaced and substituted by the amine HNR_2R_3 under reaction conditions which do not modify the other portions of the molecule. The amine reactant is usually employed in a molar excess over the activated ester in a solvent which is selected from

those mentioned above and the lower alkanols. The reaction temperature generally ranges between 0°C and 100°C. Examples of the activated ester include lower alkyl esters wherein the lower alkyl moiety is optionally substituted by cyano and nitro, phenyl esters substituted by halo and nitro groups as well as the ester moiety contained in GE 2270 factor A₂.

5 It is evident that in many instances a compound of the invention may be prepared in more than one way and that a compound of the invention may be transformed into another by means of known per se reactions.

For instance when the HNR₂R₃ amine contains a carboxy or an ester function which can be further converted into the corresponding amide derivative, a desired compound of formula I may be prepared by
10 condensing first said amine with the selected GE 2270 starting material and then converting the carboxy or ester function to amide by reaction with the appropriate amine.

The following tables list the structure formulas of some representative compounds of the invention (TABLE I) and their methods of preparation (described in details in the Experimental Section), starting materials and reaction yields (TABLE II).

15

20

25

30

35

40

45

50

55

In the following Table (TABLE I) the structure formulas of representative examples of compounds of the invention are reported.

TABLE I


COMPOUND NO.	Y	R	R ₁	R ₄
1	-NH CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
2	-NH CHCOOH CH ₂ CH ₂ CH ₂ NH ₂	CH ₂ O-CH ₃	CH ₃	CH ₃
3	-NH CHCOOH CH ₂ OH	CH ₂ O-CH ₃	CH ₃	CH ₃
4	-NH CHCOOH CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
5	-NH CH-COOH CH ₂ - 	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)

COMPOUND NO.	Y	R	R ₁	R ₄
6	$\begin{array}{c} \text{-NHCHCOOH} \\ \\ \text{CH}_2\text{CH}(\text{CH}_3)_2 \end{array}$	CH ₂ O-CH ₃	CH ₃	CH ₃
7	$\begin{array}{c} \text{-NHCH-COOH} \\ \\ \text{CH}_2-\text{C}_6\text{H}_5 \end{array}$	CH ₂ O-CH ₃	CH ₃	CH ₃
8	$\begin{array}{c} \text{-NHCHCOOH} \\ \\ \text{CH}_2\text{CH}_2\text{SCH}_3 \end{array}$	CH ₂ O-CH ₃	CH ₃	CH ₃
9	$\begin{array}{c} \text{COOH} \\ \\ \text{N} \end{array}$	CH ₂ O-CH ₃	CH ₃	CH ₃
10	$\begin{array}{c} \text{HO} \\ \\ \text{CH} \end{array}$	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)



COMPOUND NO.	Y	R	R ₁	R ₄
11	$\text{-NHCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}(\text{NH}_2)\text{COOH}$	$\text{CH}_2\text{O-CH}_3$	CH_3	CH_3
12	$\text{-NHCH}_2\text{CONHCH}(\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2)\text{COOH}$	$\text{CH}_2\text{O-CH}_3$	CH_3	CH_3
13		$\text{CH}_2\text{O-CH}_3$	CH_3	CH_3
14		$\text{CH}_2\text{O-CH}_3$	CH_3	CH_3

TABLE I (continued)

COMPOUND NO.	Y	R	R ₁	R ₄
15		CH ₂ O-CH ₃	CH ₃	CH ₃
16		CH ₂ O-CH ₃	CH ₃	CH ₃
17		CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)

COMPOUND NO.	Y	R	R ₁	R ₄
18	-NH CH ₂ CH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
19	-NH CH ₂ CH ₂ CH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
20	-NH CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
21	-NH CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
22	-NH CH ₂ CH ₂ SO ₃ H	CH ₂ O-CH ₃	CH ₃	CH ₃
23	-NH CH ₂ CH ₂ CH ₂ SO ₃ H	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)




COMPOUND NO.	Y	R	R ₁	R ₄
24	-NHCH ₂ CH ₂ CH ₂ PO ₃ H ₂	CH ₂ O-CH ₃	CH ₃	CH ₃
25	-NHCH ₂ CH ₂ CH ₂ CH ₂ CH ₂ PO ₃ H ₂	CH ₂ O-CH ₃	CH ₃	CH ₃
26	-NHCH ₂ - 	CH ₂ O-CH ₃	CH ₃	CH ₃
27	-N 	CH ₂ O-CH ₃	CH ₃	CH ₃
28	-NHCH ₂ CH ₂ CH ₂ CH ₂ - 	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)

COMPOUND NO.	Y	R	R ₁	R ₄
29	$ \begin{array}{c} \text{N-CH}_2\text{CHCHCHCHCH}_2\text{OH} \\ \quad \quad \quad \\ \text{CH}_3 \quad \text{OH} \quad \text{OH} \quad \text{OH} \end{array} $	CH ₂ O-CH ₃	CH ₃	CH ₃
30		CH ₂ O-CH ₃	CH ₃	CH ₃
31	$ \begin{array}{c} \text{O} \\ \\ \text{-NHCH}_2\text{CH}_2\text{CH}_2\text{NH-} \end{array} $	CH ₂ O-CH ₃	CH ₃	CH ₃
32	-NHCH ₂ CH ₂ N(CH ₃) ₂	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)



COMPOUND NO.	Y	R	R ₁	R ₄
33		CH ₂ O-CH ₃	CH ₃	CH ₃
34		CH ₂ O-CH ₃	CH ₃	CH ₃
35	-NH ₂	CH ₂ O-CH ₃	CH ₃	CH ₃
36	-NHCH ₂ CH ₂ CH ₂ NH ₂	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)

COMPOUND NO.	Y	R	R ₁	R ₄
37	-NH CH ₂ CHO	CH ₂ O-CH ₃	CH ₃	CH ₃
38	-NH CH ₂ CH ₂ NHCH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
39	-NH CH ₂ CH ₂ SCH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃
40	-NH CH ₂ CH ₂ CH ₂ CH = CHCOOH	CH ₂ O-CH ₃	CH ₃	CH ₃
41	-NH CH ₂ CH ₂ OCH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₃

TABLE I (continued)

COMPOUND NO.	Y	R	R ₁	R ₄
42	-NH CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	CH ₂ O-CH ₃	CH ₃	CH ₂ OH
43	-NH CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	H	H	CH ₃
44	-NH CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	CH ₂ OH	CH ₃	CH ₃

Compounds No. 2, 11, 12, 34, 36 were isolated as trifluoroacetate salts

TABLE II


COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
1	$A_3 + HCl.NH_2CH_2COOEt$	A ₁	80%
2	$A_3 + HCl.NH_2CHCOOMe$ $CH_2CH_2CH_2NH.Cbz$	A ₁	72%
3	$A_3 + HCl.NH_2CHCOOMe$ CH_2OH	A ₁	70%
4	$A_3 + HCl.NH_2CHCOOMe$ CH_2COOMe	A ₁	54%
5	$A_3 + HCl.NH_2CH-COOMe$ CH_2 	A ₁	70%
6	$A_3 + HCl.NH_2CHCOOMe$ $CH_2CH(CH_3)_2$	A ₁	70%

TABLE II (continued)





COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
7	$A_3 + HCl.NH_2CHCOOMe$ 	A ₁	60%
8	$A_3 + HCl.NH_2CHCOOMe$ 	A ₁	70%
9	$A_3 + HCl.HN$ 	A ₁	75%
10	$A_3 + HCl.NH_2CHCOOMe$ 	A ₁	70%

TABLE II (continued)

COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
11	$A_3 + NH_2CH_2CH_2CH_2CH_2CH_2COOH$ $NH.Cbz$	B ₁	64%
12	$A_3 + TFA.NH_2CH_2CONHCH_2COOH$ $CH_2CH_2CH_2CH_2NH.Cbz$	B ₁	74%
13	 $\underline{3} + HCl.HN$	C ₁	70%
14	 $\underline{1} +$	C	83%

TABLE II (continued)








COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
15	$A_3 + HCl.NH_2CHCON$  	A	60%
	or $\underline{5} +$ 	C	70%
16	$\underline{10} +$ 	C	60%
17	$\underline{6} +$ 	C	65%

TABLE II (continued)

COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
18	A ₃ + HCl.NH ₂ CH ₂ CH ₂ CH ₂ COOMe	A ₁	73%
19	A ₃ + HCl.NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOMe	A ₁	77%
	A ₃ + NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	B	70%
20	A ₃ + PTSA.NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOMe	A ₁	75%
21	A ₃ + PTSA.NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOMe	A ₁	70%
22	A ₃ + NH ₂ CH ₂ CH ₂ SO ₃ H	B	20%

TABLE II (continued)

COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
23	$A_3 + NH_2CH_2CH_2CH_2SO_3H$	B	25%
24	$A_3 + NH_2CH_2CH_2CH_2PO_3H_2$	B	40%
25	$A_3 + NH_2CH_2CH_2CH_2CH_2CH_2PO_3H_2$	B	35%
26	$A_3 + NH_2CH_2$ 	B	60%
27	$A_3 + HN$ 	B	50%

COMPOUND	STARTING MATERIALS	METHOD	OVERALL YIELD
1	2	3	4




COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
28	$A_3 + NH_2CH_2CH_2CH_2CH_2CH_2 - \text{C}_5\text{H}_3\text{N}_3$ 	B	65%
29	$A_3 + NH-CH_2CH(CH_3)CH(OH)CH(OH)CH_2OH$	A	20%
30	$A_3 + \text{Cyclohexane-1,2,3-triol-4-amine.HCl}$ 	A	80%
31	$36 + \text{2,5-bis(4-phenylthioxy)phenyl 1,3-dithiolane-2-carboxylate}$ 	C ₁	80%

TABLE II (continued)



COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
32	$A_3 + NH_2CH_2CH_2N(CH_3)_2$	A	75%
33	$A_3 + NH_2$ 	A	60%
34	$A_3 + NH$ 	A ₁	50%
35	$A_2 + NH_3$ in MeOH	D	83%
36	$A_3 + NH_2CH_2CH_2CH_2NH.Boc$	A ₁	70%

TABLE II (continued)

COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
37	$ \begin{array}{c} \text{OCH}_3 \\ \diagup \\ \text{A}_3 + \text{NH}_2\text{CH}_2\text{CH} \\ \diagdown \\ \text{OCH}_3 \end{array} $	A ₁	65%
38	$\underline{37} + \text{HCl.NH}_2\text{CH}_2\text{CH}_2\text{COOCH}_2\text{CH}_3$	C ₁	20%
39	$\text{A}_3 + \text{TFA.NH}_2\text{CH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{COOCH}_3$	A ₁	33%
40	$\text{A}_3 + \text{TFA.NH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH-COOH}$	B	51%

TABLE II (continued)

COMPOUND NO.	STARTING MATERIALS (GE 2270 FACTOR + AMINE REACTANT)	METHOD	OVERALL YIELD
41	A ₃ + TFA.NH ₂ CH ₂ CH ₂ OCH ₂ CH ₂ COOH	B	37%
42	C _{2a} + HCl.NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOCH ₃	F	40%
	NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	G	35%
43	D ₁ + HCl.NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOCH ₃	H	50%
	NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	I	40%
44	D ₂ + HCl.NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOCH ₃	J	35%
	NH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ COOH	K	30%

TFA = trifluoroacetic acid
PTSA = p-toluenesulfonic acid

HPLC Analysis

The following table (TABLE III) reports the R_f of representative examples of compounds of this invention.

Analysis were run with a Varian model 5000 LC pump equipped with a 10 μ l loop injector and a Varian 2050 variable wavelength detector at 254 nm.

Columns: Pre-column LiChroCart-LiChrosorb RP-8 (5 μ m) followed by a column LiChroCart 125-4 LiChrospher 100 RP-8 (5 μ m)

5 Eluents: A 0.05 M aq. HCOONH₄
 B CH₃CN
 C THF

Method A: isocratic 44% of B in A

 Flow rate: 0.7 ml/min

10 Method B: isocratic 40% of B in A

 Flow rate: 0.7 ml/min

Method C: isocratic 38% of B in A

 Flow rate: 0.5 ml/min

Method D: isocratic 30% of B in A

15 Flow rate: 0.7 ml/min

Method E: isocratic 38% of B in A Flow rate: 0.7 ml/min

Method F: gradient from 38 to 55% of B in A in 11 min according to the following program

Time (min)	%B in A
0	38
6	38
7	45
10	45
11	55

25 Flow rate: 0.7 ml/min

Method G: gradient from 38 to 55% of B in A in 25 min according to the following program

Time (min)	%B in A
0	38
6	38
10	44
15	44
25	55

30 Flow rate: 0.7 ml/min

Method H: isocratic 55% of B in A

 Flow rate: 0.7 ml/min

Method I: isocratic 60% of B in A

 Flow rate: 0.7 ml/min

Method L: isocratic 48% of B in A

45 Flow rate: 0.7 ml/min

Method M: gradient according to the following program:

50

55

Time (min)	%A	%B	%C
0	74	10	16
20	62	19	19

Flow rate: 0.7 ml/min

TABLE III
HPLC Analysis

Compound No.	Method	R _t (min)	K
1	A	2.56	0.92
2	B	4.09	1.15
3	C	6.21	1.12
4	D	14.70	1.79
5	E	5.99	1.28
6	A	4.05	1.46
7	A	4.52	1.63
8	A	3.44	1.24
9	E	5.32	1.18
10	E	4.22	0.90
11	F	14.30	3.05
12	F	5.99	1.28
13	E	4.88	1.04
14	G	14.09	3.01
15	G	17.60	3.76
16	G	13.77	2.94
17	G	23.75	5.07
18	G	7.49	1.60

K = Relative Retention time

TABLE III (continued)
HPLC Analysis

Compound No.	Method	R _t (min)	K
19	F	8.84	1.89
20	G	17.77	3.78
21	G	31.10	6.64
22	F	5.01	1.07
23	F	4.40	0.94
24	F	6.27	1.34
25	F	11.17	2.39
26	F	29.04	6.02
27	F	6.28	1.34
28	F	12.14	2.59
29	F	8.02	1.71
30	E	6.81-7.61 anomeric mixture	1.45-1.62 anomeric mixture
31	F	17.64	3.76
32	B	9.64	2.70
33	H	15.10	7.40
34	I	7.22	3.92
35	L	10.28	4.11
36	F	19.32	4.13

K = Relative Retention time

TABLE III (continued)
HPLC Analysis

Compound No.	Method	R _t (min)	K
37	F	14.56	3.11
38	F	11.00	2.35
39	F	9.48	2.02
40	F	6.84	1.46
41	F	3.95	0.84
42	M	17.23	1.32
43	M	15.76	1.52
44	M	16.64	1.50
19	M	20.81	1.32

K = Relative Retention time

K = Relative Retention time =

$R_t \text{ amide} / R_t \text{ GE 2270 proper starting material (i.e. the compound of formula II wherein W is COOH)}$

EXPERIMENTAL SECTION

TABLE IV - N.M.R.

The ¹H-NMR spectra were recorded with a Bruker spectrometer in DMSO-d₆ - (hexadeuterodimethylsulfoxide) using TMS as the internal standard (δ 0.00 ppm) [δ, ppm, multiplicity] at 250 MHz and/or 500 MHz (s = singlet, br s = broad singlet, d = doublet, dd = doublet of doublets, t = triplet, m = multiplet)

TABLE V - I.R.

The infrared spectra (IR) were recorded with a Perkin Elmer mod. 580 spectrophotometer in nujol mull.

TABLE VI - U.V.

The ultraviolet absorption spectra were recorded with a Perkin Elmer Model 320 spectrometer.

It will be clear to the skilled technician that the data represented in TABLES IV, V and VI below, do not represent all the values of the peaks obtained but only the values of those peaks which permit to characterize the single substance.

TABLE IV - N.M.R. Spectra

COMPOUND NO.	¹ H-NMR (DMSOd ₆) δ(ppm)
1	0.84 (d, 3H); 0.87 (d, 3H); 2.57 (s, 3H); 3.39 (s, 3H); 3.77 (dd, 1H); 3.99 (d, 2H); 4.25 (dd, 1H); 4.96 (s, 2H); 7.36-7.22 (m, 7H); 8.28 (s, 1H); 8.50 (s, 1H); 8.59 (s, 1H)
2	0.79 (d, 3H); 0.85 (d, 3H); 2.05-1.70 (m, 4H); 2.54 (s, 3H); 3.33 (s, 3H); 3.65 (m, 2H); 3.81 (dd, 1H); 4.10 (m, 1H); 4.35 (dd, 1H); 4.99 (s, 2H); 7.35-7.05 (m, 7H); 8.20 (s, 1H); 8.42 (s, 1H); 8.58 (s, 1H)
3	0.84 (d, 3H); 0.87 (d, 3H); 2.58 (s, 3H); 3.37 (s, 3H); 3.80 (dd, 2H); 3.84 (dd, 1H); 3.91 (dd, 1H); 4.26 (dd, 1H); 4.55 (m, 1H); 4.97 (s, 2H); 7.36-7.20 (m, 7H); 8.29 (s, 1H); 8.55 (s, 1H); 8.59 (s, 1H)
4	0.85 (d, 3H); 0.89 (d, 3H); 2.58 (s, 3H); 2.90 (m, 2H); 3.38 (s, 3H); 3.70 (dd, 1H); 4.29 (dd, 1H); 4.85 (m, 1H); 4.98 (s, 2H); 7.40-7.20 (m, 7H); 8.28 (s, 1H); 8.52 (s, 1H); 8.58 (s, 1H)
5	0.85 (d, 3H); 0.88 (d, 3H); 2.58 (s, 3H); 3.11 (m, 2H); 3.26 (br, s, 1H); 3.38 (s, 3H); 3.78 (dd, 1H); 4.28 (dd, 1H) 4.64 (m, 1H); 4.97 (s, 2H); 6.68 (d, 1H); 7.09 (d, 1H); 7.40-7.20 (m, 7H); 8.27 (s, 1H); 8.47 (s, 1H); 8.59 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
6	0.84 (d, 3H); 0.87 (d, 3H); 0.92 (d, 3H); 0.95 (d, 3H); 1.69 (m, 2H); 1.86 (m, 1H); 2.57 (s, 3H); 3.37 (s, 3H); 3.78 (dd, 1H); 4.26 (dd, 1H); 4.53 (m, 1H); 4.97 (s, 2H); 7.38-7.20 (m, 7H); 8.28 (s, 1H); 8.46 (s, 1H); 8.59 (s, 1H)
7	0.84 (d, 3H); 0.88 (d, 3H); 2.58 (s, 3H); 3.20 (m, 2H); 3.37 (s, 3H); 3.77 (dd, 1H); 4.25 (dd, 1H); 4.73 (m, 1H); 7.40-7.2 (m, 12H); 8.28 (s, 1H); 8.47 (s, 1H) 8.59 (s, 1H)
8	0.85 (d, 3H); 0.89 (d, 3H); 2.08 (s, 3H); 2.16 (m, 2H); 2.56 (m, 2H); 2.57 (s, 3H); 3.40 (s, 3H); 3.79 (dd, 1H); 4.27 (dd, 1H); 4.61 (m, 1H); 5.00 (s, 2H); 7.37-7.20 (m, 7H); 8.29 (s, 1H); 8.52 (s, 1H); 8.60 (s, 1H)
9	0.84 (d, 3H); 0.88 (d, 3H); 2.45-1.70 (m, 4H); 2.58 (s, 3H); 3.37 (s, 3H); 3.68 (m, 2H); 3.78 (dd, 1H); 4.10 (m, 1H); 4.27 (dd, 1H); 4.49 (m, 1H); 7.35-7.22 (m, 7H); 8.27 (s, 1H); 8.50 (s, 1H); 8.59 (s, 1H)
10	0.85 (d, 3H); 0.88 (d, 3H); 1.19 (d, 3H); 2.59 (s, 3H); 3.39 (s, 3H); 3.78 (dd, 1H); 4.30 (m, 2H); 4.48 (dd, 1H); 4.99 (s, 2H); 7.4-7.2 (m, 7H); 8.33 (s, 1H); 8.49 (s, 1H); 8.60 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
11	0.84 (d, 3H); 0.88 (d, 3H); 1.55-1.35 (m, 2H); 1.61 (m, 2H); 1.83 (m, 2H); 2.58 (s, 3H); 3.34 (m, 2H); 3.38 (s, 3H); 3.79 (dd, 1H); 3.91 (br; s, 1H); 4.29 (dd, 1H); 4.97 (s, 2H); 7.35-7.13 (m, 7H); 8.27 (s, 1H); 8.43 (s, 1H); 8.59 (s, 1H)
12	0.85 (d, 3H); 0.88 (d, 3H); 1.37 (m, 2H); 1.70-1.49 (m, 3H); 1.75 (m, 1H); 2.58 (s, 3H); 2.76 (m, 2H); 3.38 (s, 3H); 3.78 (dd, 1H); 4.03 (m, 2H); 4.28 (m, 2H); 4.97 (s, 2H); 7.35-7.20 (m, 7H); 8.28 (s, 1H); 8.49 (s, 1H); 8.59 (s, 1H)
13	0.84 (d, 3H); 0.88 (d, 3H); 1.98-1.82 (m, 2H); 2.18 (m, 2H); 2.56 (s, 3H); 2.69 (dd, 2H); 3.36 (s, 3H); 3.85-3.62 (m, 3H); 4.31 (m, 2H); 4.85 (m, 1H); 4.96 (s, 2H); 7.38-7.19 (m, 7H); 8.24 (s, 1H); 8.55 (s, 1H); 8.63 (s, 1H)
14	0.85 (d, 3H); 0.88 (d, 3H); 1.99-1.82 (m, 3H); 2.06 (m, 1H); 2.58 (s, 3H); 3.58 (m, 1H); 3.67 (m, 1H); 3.79 (dd, 1H); 4.18 (d, 2H); 4.28 (dd, 1H); 4.97 (s, 2H); 6.93 (s, 1H); 7.36-7.28 (m, 8H); 8.28 (s, 1H); 8.52 (s, 1H); 8.59 (s, 1H)
15	0.85 (d, 3H); 0.88 (d, 3H); 2.07-1.63 (m, 4H); 2.58 (s, 3H); 2.99 (dd, 1H); 3.09 (dd, 1H); 3.38 (s, 3H); 3.51 (m, 1H); 3.77 (m, 2H); 4.30 (m, 2H); 4.89 (m, 1H); 4.98 (s, 2H); 6.66 (d, 1H); 6.95 (br; s, 1H); 7.16 (d, 1H); 7.39-7.20 (m, 8H); 8.23 (s, 1H); 8.42 (s, 1H); 8.58 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ (ppm)
16	0.84 (d, 3H); 0.87 (d, 3H); 1.20 (d, 3H); 1.98-1.80 (m, 3H); 2.08 (m, 1H); 2.56 (s, 3H); 3.36 (s, 3H); 3.85-3.71 (m, 2H); 4.13 (m, 1H); 4.28 (dd, 1H); 4.31 (dd, 1H); 4.73 (m, 1H); 5.05 (d, 1H); 6.89 (br; s, 1H); 7.15 (br; s, 1H); 7.38-7.19 (m, 7H); 8.26 (s, 1H); 8.51 (s, 1H); 8.56 (s, 1H)
17	0.84 (d, 3H); 0.87 (d, 3H); 0.94 (d, 3H); 0.98 (d, 3H); 2.10-1.62 (m, 7H); 2.56 (s, 3H); 3.36 (s, 3H); 3.65 (m, 1H); 3.88-3.70 (m, 2H); 4.31 (m, 2H); 4.88 (m, 1H); 4.96 (s, 2H); 6.79 (br, s, 1H); 7.18 (br; s, 1H); 7.35-7.20 (m, 7H); 8.25 (s, 1H); 8.48 (s, 1H); 8.56 (s, 1H)
18	0.84 (d, 3H); 0.88 (d, 3H); 1.81 (m, 2H); 2.30 (t, 2H); 2.58 (s, 3H); 3.35 (m, 2H); 3.37 (s, 3H); 3.78 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 2H); 7.35-7.20 (m, 7H); 8.27 (s, 1H); 8.46 (s, 1H); 8.59 (s, 1H)
19	0.84 (d, 3H); 0.87 (d, 3H); 1.35 (m, 2H); 1.56 (m, 4H); 2.22 (t, 2H); 2.58 (s, 3H); 3.36 (m, 2H); 3.38 (s, 3H); 3.80 (dd, 1H); 4.29 (dd, 1H); 4.97 (s, 2H); 7.42-7.22 (m, 7H); 8.29 (s, 1H); 8.45 (s, 1H); 8.62 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSOd ₆) δ(ppm)
20	0.84 (d, 3H); 0.88 (d, 3H); 1.31 (br; s, 6H); 1.51 (m, 2H); 1.57 (m, 2H); 2.19 (t, 2H); 2.58 (s, 3H); 3.32 (m, 2H); 3.37 (s, 3H); 3.79 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 2H); 7.38-7.19 (m, 7H); 8.27 (s, 1H); 8.45 (s, 1H); 8.59 (s, 1H)
21	0.84 (d, 3H); 0.88 (d, 3H); 1.41-1.20 (m, 12H); 1.47 (m, 2H); 1.57 (m, 2H); 2.17 (t, 2H); 2.58 (s, 3H); 3.29 (m, 2H); 3.38 (s, 3H); 3.79 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 2H); 7.38-7.19 (m, 7H); 8.27 (s, 1H); 8.43 (s, 1H); 8.59 (s, 1H)
22	0.85 (d, 3H); 0.87 (d, 3H); 2.57 (s, 3H); 2.79 (t, 2H); 3.37 (s, 3H); 3.59 (t, 2H); 3.78 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 1H); 7.41-7.20 (m, 7H); 8.27 (s, 1H); 8.44 (s, 1H); 8.57 (s, 1H)
23	0.84 (d, 3H); 0.87 (d, 3H); 1.67 (m, 2H); 2.53 (t, 2H); 2.57 (s, 3H); 3.26 (t, 2H); 3.37 (s, 3H); 3.78 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 2H); 7.41-7.26 (m, 7H); 8.26 (s, 1H); 8.44 (s, 1H); 8.57 (s, 1H)
24	0.85 (d, 3H); 0.88 (d, 3H); 1.58 (m, 2H); 1.79 (m, 2H); 2.58 (s, 3H); 3.38 (s, 3H); 3.50 (m, 2H); 3.78 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 2H); 7.38-7.21 (m, 7H); 8.27 (s, 1H); 8.45 (s, 1H); 8.59 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
25	0.84 (d, 3H); 0.88 (d, 3H); 1.65-1.35 (m, 8H); 4.58 (s, 3H); 3.38 (s, 3H); 3.78 (dd, 1H); 4.28 (dd, 1H); 4.97 (s, 2H); 7.40-7.20 (m, 7H); 8.28 (s, 1H); 8.43 (s, 1H); 8.59 (s, 1H)
26	0.85 (d, 3H); 0.89 (d, 3H); 2.56 (s, 3H); 3.36 (s, 3H); 3.80 (dd, 1H); 4.31 (dd, 1H); 4.62 (br, s, 2H); 4.96 (s, 2H); 7.39-7.15 (m, 7H); 7.47 (d, 2H); 7.90 (d, 2H); 8.26 (s, 1H); 8.41 (s, 1H); 8.58 (s, 1H)
27	0.84 (d, 3H); 0.88 (d, 3H); 1.62 (br, s, 2H); 1.92 (br, s, 2H); 2.58 (s, 3H); 2.60 (m, 1H); 3.38 (s, 3H); 3.79 (dd, 1H); 4.16 (m, 2H); 4.29 (dd, 1H); 4.38 (m, 2H); 7.35-7.19 (m, 7H); 8.25 (s, 1H); 8.29 (s, 1H); 8.57 (s, 1H)
28	0.85 (d, 3H); 0.89 (d, 3H); 1.39 (m, 2H); 1.61 (m, 2H); 1.76 (m, 2H); 2.58 (s, 3H); 2.88 (t, 2H); 3.33 (m, 2H); 3.80 (dd, 1H); 4.29 (dd, 1H); 4.98 (s, 2H); 7.34-7.20 (m, 7H); 8.26 (s, 1H); 8.45 (s, 1H); 8.58 (s, 1H)
29	0.84 (d, 3H); 0.88 (d, 3H); 2.58 (s, 6H); 3.38 (s, 3H); 3.70-3.41 (m, 5H); 3.89-3.75 (m, 2H); 3.98 (br, s, 1H); 4.35-4.26 (m, 2H); 4.97 (s, 2H); 7.35-7.21 (m, 7H); 8.26 (s, 1H); 8.28 (s, 1H); 8.58 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
30	0.84 (d, 3H); 0.88 (d, 3H); 2.58 (s, 3H); 3.29-3.14 (m, 2H); 3.38 (s, 3H); 3.90-3.49 (m, 4H); 4.29 (dd, 1H); 4.92 (m, 1H); 4.97 (s, 2H); 5.12 (t, 1H); 7.35-7.18 (m, 7H); 8.26 (s, 1H); 8.51 (s, 1H); 8.58 (s, 1H)
31	0.86 (d, 3H); 0.89 (d, 3H); 1.81 (m, 2H); 2.59 (s, 3H); 3.32 (m, 4H); 3.39 (s, 3H); 3.80 (dd, 1H); 4.30 (dd, 1H); 4.99 (s, 2H); 6.75 (d, 1H); 7.41-7.18 (m, 9H); 8.28 (s, 1H); 8.46 (s, 1H); 8.59 (s, 1H)
32	0.85 (d, 3H); 0.88 (d, 3H); 2.21 (s, 6H); 2.59 (s, 3H); 3.38 (s, 3H); 3.43 (m, 4H); 3.81 (dd, 1H); 4.31 (dd, 1H); 4.98 (s, 2H); 7.45-7.19 (m, 7H); 8.28 (s, 1H); 8.45 (s, 1H); 8.61 (s, 1H)
33	0.86 (d, 3H); 0.90 (d, 3H); 1.91-1.70 (m, 2H); 2.26-2.05 (m, 2H); 2.60 (s, 3H); 2.91-2.69 (m, 4H); 3.40 (s, 3H); 3.51 (br; s, 2H); 3.95-3.75 (m, 2H); 4.30 (dd, 1H); 4.99 (s, 2H); 7.41-7.18 (m, 12H); 8.28 (s, 1H); 8.45 (s, 1H); 8.66 (s, 1H)
34	0.85 (d, 3H); 0.89 (d, 3H); 1.81-1.49 (m, 4H); 2.01-1.88 (m, 2H); 2.59 (s, 3H); 2.98-2.65 (m, 4H); 3.39 (s, 3H); 3.80-3.51 (m, 4H); 3.81 (dd, 1H); 4.31 (dd, 1H); 4.99 (s, 2H); 7.41-7.18 (m, 7H); 7.90-7.65 (m, 6H); 8.25 (s, 1H); 8.36 (s, 1H); 8.61 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
35	0.85 (d, 3H); 0.88 (d, 3H); 2.59 (s, 3H); 3.39 (s, 3H); 3.79 (dd, 1H); 4.29 (dd, 1H); 4.98 (s, 2H); 7.40-7.19 (m, 7H); 7.72 (br, s, 1H); 8.03 (br, s, 1H); 8.28 (s, 1H); 8.47 (s, 1H); 8.60 (s, 1H)
36	0.85 (d, 3H); 0.88 (d, 3H); 1.87 (m, 2H); 2.54 (s, 3H); 2.89 (m, 2H); 3.37 (s, 3H); 3.42 (m, 2H); 3.79 (dd, 1H); 4.29 (dd, 1H); 4.98 (s, 2H); 7.38-7.20 (m, 7H); 7.69 (br, s, 3H); 8.29 (s, 1H); 8.49 (s, 1H); 8.61 (s, 1H)
37	0.83 (d, 3H); 0.87 (d, 3H); 1.32 (m, 1H); 2.16 (m, 1H); 2.46 (d, 3H); 2.57 (s, 3H); 2.71 (m, 1H); 3.37 (s, 3H); 3.78 (dd, 1H); 4.16 (d, 1H); 4.26 (dd, 1H); 4.67 (m, 1H); 4.96 (s, 2H); 6.02 (d, 1H); 6.35 (dd, 1H); 7.35-7.20 (m, 7H); 8.28 (s, 1H); 8.49 (s, 1H); 8.60 (s, 1H); 9.61 (s, 1H)
38	0.83 (d, 3H); 0.87 (d, 3H); 1.25 (m, 1H); 2.2 (m, 1H); 2.5 (s, 3H); 2.70 (m, 3H); 3.35 (s, 3H); 3.63 (m, 1H); 3.79 (d, 1H); 4.27 (dd, 1H); 4.97 (s, 2H); 7.4-7.15 (m, 7H); 8.28 (s, 1H); 8.53 (s, 1H); 8.61 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
39	0.83 (d, 3H); 0.87 (d, 3H); 1.32 (m, 1H); 2.16 (m, 1H); 2.47 (d, 3H); 2.57 (s, 3H); 2.72 (m, 4H); 3.37 (s, 3H); 3.50 (m, 2H); 3.78 (dd, 1H); 4.27 (dd, 1H); 4.97 (s, 2H); 7.40-7.20 (m, 7H); 8.28 (s, 1H); 8.49 (s, 1H); 8.60 (s, 1H)
40	0.83 (d, 3H); 0.87 (d, 3H); 1.32 (m, 1H); 1.71 (m, 2H); 2.25-2.14 (m, 3H); 2.46 (d, 3H); 2.57 (s, 3H); 2.7 (m, 1H); 3.37 (s, 3H); 3.76 (dd, 1H); 4.27 (dd, 1H); 4.97 (s, 2H); 5.81 (d, 1H, J=15.7 Hz); 6.78 (m, 1H); 7.39-7.12 (m, 7H); 8.28 (s, 1H); 8.45 (s, 1H); 8.60 (s, 1H)
41	0.86 (d, 3H); 0.89 (d, 3H); 1.43 (m, 1H); 2.19 (m, 1H); 2.47 (d, 3H); 2.59 (s, 3H); 2.72 (m, 1H); 3.39 (s, 3H); 3.50 (t, 2H); 3.58 (t, 2H); 3.68 (t, 2H); 3.79 (dd, 1H); 4.99 (s, 2H); 7.42-7.20 (m, 7H); 8.27 (s, 1H); 8.47 (s, 1H); 8.59 (s, 1H)
42	0.83 (d, 3H); 0.85 (d, 3H); 1.2-1.4 (m, 3H); 1.5-1.65 (m, 4H); 2.22 (t, 3H); 2.60 (d, 1H); 2.69 (d, 1H); 3.37 (s, 3H); 3.79 (dd, 1H); 4.27 (dd, 1H); 4.86 (d, 2H); 4.97 (s, 2H); 5.00 (dd, 1H); 5.1-5.4 (m, 3H); 5.74 (t, 1H); 6.00 (d, 1H); 7.2-7.4 (m, 7H); 8.27 (s, 1H); 8.44 (s, 1H); 8.62 (s, 1H)

TABLE IV - N.M.R. Spectra (continued)

COMPOUND NO.	¹ H-NMR (DMSO-d ₆) δ(ppm)
43	0.84 (d, 3H); 0.89 (d, 3H); 1.4-1.2 (m, 3H); 1.65-1.50 (m, 4H); 2.23 (t, 3H); 2.59 (s, 3H); 2.79 (m, 1H); 3.87 (m, 1H); 4.25 (m, 1H); 5.04 (t, 1H); 5.35-5.20 (m, 3H); 6.09 (d, 1H); 6.67 (br, s, 1H); 7.04 (br, s, 1H); 7.35-7.15 (m, 6H); 8.24 (s, 1H); 8.26 (s, 1H); 8.45 (s, 1H); 8.61 (s, 1H)
44	0.84 (d, 3H); 0.88 (d, 3H); 1.4-1.25 (m, 3H); 1.65-1.50 (m, 4H); 2.23 (t, 3H); 2.58 (s, 3H); 2.75 (m, 1H); 3.78 (dd, 1H); 4.28 (dd, 1H); 4.98 (m, 3H); 5.35-5.15 (m, 3H); 6.03 (m, 2H); 7.42-7.15 (m, 7H); 8.30 (s, 1H); 8.45 (s, 1H); 8.62 (s, 1H)

TABLE V - I.R. Spectra

COMPOUND NO.	I.R. (nujol cm ⁻¹)
1	3370; 3110; 1730; 1655; 1545; 1520
2	3350; 3110; 1720; 1650; 1535; 1500
3	3340; 3105; 1720; 1645; 1535; 1500
4	3360; 1725; 1640; 1535
5	3350; 3110; 1725; 1650; 1535; 1510
6	3370; 3105; 1725; 1655; 1535; 1500
7	3360; 3100; 1725; 1655; 1535; 1490
8	3370; 3105; 1725; 1655; 1535; 1505
9	3370; 3110; 3100; 1725; 1657; 1550; 1530; 1505
10	3370; 3105; 1730; 1655; 1540; 1510
11	3359; 3115; 1653; 1551; 1510;
12	3360; 3113; 1720; 1662; 1547; 1510

TABLE V - I.R. Spectra (continued)

COMPOUND NO.	I.R. (nujol cm ⁻¹)
13	3370; 3110; 1720; 1655; 1530; 1505
14	3350; 3120; 1655; 1535; 1500
15	3350; 3100; 1650; 1530; 1510;
16	3340; 3105; 1650; 1530
17	3340; 3100; 1655; 1530
18	3350; 3100; 1710; 1645; 1540
19	3360; 3115; 1720; 1665; 1540; 1506;
20	3350; 3113; 1720; 1659; 1549; 1506
21	3340; 1710; 1645; 1540; 1500
22	3304; 1653; 1540
23	3333; 1657; 1547; 1092; 1038;
24	3354; 3113; 1653; 1550; 1506; 1245

TABLE V - I.R. Spectra (continued)

COMPOUND NO.	I.R. (nujol cm ⁻¹)
25	3348; 3111; 1660; 1548; 1507; 1245
26	3315; 1653; 1539; 1238
27	3361; 3113; 1720; 1653; 1531; 1507; 1092
28	3333; 1653; 1547; 1494; 1243
29	3356; 3114; 1653; 1508; 1088
30	3360; 1670; 1505; 1200
31	3351; 3115; 1653; 1549; 1509; 1250
32	3370; 3110; 1655; 1545; 1500; 1245
33	3350; 1655; 1530; 1490; 1220
34	3360; 3105; 1650; 1545; 1510; 1240
35	3320; 1747; 1650; 1540; 1225
36	3330; 1662; 1547; 1496; 1201

TABLE V - I.R. Spectra (continued)

COMPOUND NO.	I.R. (nujol cm ⁻¹)
37	3327; 1730; 1653; 1464; 1377
38	3355; 1720; 1657; 1543; 1377
39	3321; 1717; 1652; 1545
40	3337; 1665; 1549
41	3341; 1721; 1653; 1548; 1377
42	3335; 1722; 1647; 1543
43	3317; 1665; 1539
44	3317; 1720; 1649; 1545

TABLE VI
U.V. DATA λ_{\max} ($E^{1\%}_{1\text{cm}}$)

Compound No.	MeOH	HCl 0.1N	Phosphate Buffer pH 7.38	KOH 0.1N
1	309 (290.9)	312	309 (247.9)	309 (252.7)
2	309 (257.5)	310 (222.6)	311	309 (226.3)
3	309 (297.5)	312	309 (229.6)	309 (235.8)
4	309 (245.1)	312	308 (234.7)	308 (234.1)
5	308 (173.8)	312	309 (150.0)	305 (181.2)
6	309 (277.1)	313	309 (229.8)	309 (236.7)
7	309 (258.6)	313	309 (207.9)	309 (218.9)
8	309 (279.8)	311	309 (225.9)	309 (229.4)
9	309 (261.9)	313	308 (228.1)	309 (235.0)
10	309 (279.3)	314	309 (241.8)	309 (251.1)
11	309 (216.8)	310 (178.5)	312	310 (194.9)
12	309 (226.2)	310 (188.3)	311	309 (202.4)

TABLE VI (continued)
U.V. DATA λ_{max} (E 1%)
1cm

Compound No.	MeOH	HCl 0.1N	Phosphate Buffer pH 7.38	KOH 0.1N
13	308 (237.9)	314	308 (247.4)	308 (260.3)
14	309 (263.4)	313	313	314
15	309 (222.6)	313	314	304 (169.8)
16	309 (235.6)	313	312	312
17	309 (230.3)	312	312	312
18	309 (288.8)	313	309 (239.4)	309 (248.0)
19	309 (283.2)	312	309 (220.3)	309 (230.1)
20	309	314	309	309
21	309 (271.6)	313	311 (221.6)	309 (221.6)
22	309 (190.9)	309 (152.2)	308 (160.4)	309 (165.6)
23	309 (242.2)	310 (182.2)	309 (200.9)	309 (200.9)
24	309	312	310	309

TABLE VI (continued)
U.V. DATA λ_{\max} ($E^{1\%}_{1\text{cm}}$)

Compound No.	MeOH	HCl 0.1N	Phosphate Buffer pH 7.38	KOH 0.1N
25	309	312	310	309
26	309 (260.0)	313	310 (197.7)	310 (208.6)
27	310 (264.6)	313	310 (227.4)	310 (232.1)
28	309 (260.5)	314	310 (186.8)	310 (203.6)
29	309 (243.4)	311	312	311
30	309 (248.5)	311	311	309
31	305 (253.7)	310	310	313 (249.5)
32	309 (267.9)	310 (234.9)	312	312
33	309 (247.5)	311 (234.4)	314	312
34	310 (224.0)	309 (198.2)	310	312
35	308 (269.8)	314	313	313
36	309 (243.9)	309 (205.5)	312	313

TABLE VI (continued)
U.V. DATA λ_{max} ($E^{1\%}_{1cm}$)

Compound No.	MeOH	HCl 0.1N	Phosphate Buffer pH 7.38	KOH 0.1N
37	309 (255.1)	312	314	312
38	308	308	308	308
39	308 (247.9)	312	308 (201.3)	308 (215)
40	309 (304.3)	312	309 (235.9)	309 (262.0)
41	309 (256.4)	312	309 (215.1)	309 (228.6)
42	309	312	309	307
43	309 (253.6)	313	309 (208.4)	309 (235.4)
44	309 (264.9)	314	309 (208.1)	309 (223.7)

The antimicrobial activity of the compounds of the invention can be demonstrated by a series of standard tests in vitro.

MIC for Propionibacterium acnes, and Bacteroides fragilis are determined by agar dilution (inocula $10^4/10^5$ CFU/spot). MIC for other organisms are determined by microbroth dilution (inocula 10^4 to 10^5 CFU/ml). Incubation times are 18-24 h, except for Haemophilus influenzae, P. acnes, B. fragilis (48 h). All organisms are incubated at 37°C ; H. influenzae is incubated in a 5% CO_2 atmosphere, anaerobes in an anaerobic gas mixture. Media used are: Iso-Sensitest broth (Oxoid) (Staphylococci, Streptococcus faecalis, Escherichia coli, Proteus vulgaris); brain heart infusion broth (Difco) + 1% Supplement C (Difco) (H. influenzae).

The minimal inhibitory concentrations (MIC, microgram/ml) for some microorganisms are reported below in Table VII.

TABLE VII - (MIC, microgram/ml)

STRAIN	COMPOUND NO.							
	1	2	6	7	8			
<u>Staph. aureus</u> L165 Tour	0.5	0.13	<0.13	0.25	0.25			
<u>Staph. epidermidis</u> L147 ATCC 12228	1	0.25	1	0.5	2			
<u>Staph. haemolyticus</u> L602	4	16	1	4	2			
<u>Strep. pneumoniae</u> L44 UC41	8	>128	2	4	4			
<u>Strep. faecalis</u> L149 ATCC 7080	0.25	0.06	<0.13	<0.13	0.25			
<u>Prop. acnes</u> L1014 ATCC 6919	<0.13	0.06	<0.13	<0.13	<0.13			
<u>Bact. fragilis</u> L1010 ATCC 23745	8	>128	>128	>128	32			
<u>Haemophilus influenzae</u> type B	8	>128	32	128	64			
<u>Esch. coli</u> L47 SKF 12140	>128	>128	>128	>128	>128			
<u>Prot. vulgaris</u> ATCC 881	>128	>128	>128	>128	>128			

TABLE VII - (MIC, microgram/ml) (continued)

STRAIN	COMPOUND NO.				
	14	18	19	20	26
<u>Staph. aureus</u> L165 Tour	0.06	0.25	0.06	0.25	0.25
<u>Staph. epidermidis</u> L147 ATCC 12228	0.13	0.25	0.06	0.25	0.25
<u>Staph. haemolyticus</u> L602	0.25	1	0.25	0.25	0.5
<u>Strep. pneumoniae</u> L44 UC41	> 128	1	0.25	2	2
<u>Strep. faecalis</u> L149 ATCC 7080	0.06	0.13	0.06	< 0.13	0.13
<u>Prop. acnes</u> L1014 ATCC 6919	0.03	0.016	0.008	0.03	0.008
<u>Bact. fragilis</u> L1010 ATCC 23745	> 128	2	4	> 128	32
<u>Haemophilus influenzae</u> type B	> 128	2	2	> 128	8
<u>Esch. coli</u> L47 SKF 12140	> 128	> 128	> 128	> 128	> 128
<u>Prot. vulgaris</u> ATCC 881	> 128	> 128	> 128	> 128	> 128

TABLE VII - (MIC, microgram/ml) (continued)

STRAIN	COMPOUND NO.			
	27	28	32	35
<u>Staph. aureus</u> L165 Tour	0.25	0.13	0.5	0.13
<u>Staph. epidermidis</u> L147 ATCC 12228	0.5	0.5	0.5	0.13
<u>Staph. haemolyticus</u> L602	1	1	0.5	0.5
<u>Strep. pneumoniae</u> L44 UC41	8	2	1	> 128
<u>Strep. faecalis</u> L149 ATCC 7080	1	0.06	0.25	0.06
<u>Prop. acne</u> L1014 ATCC 6919	0.03	0.008	0.13	0.004
<u>Bact. fragilis</u> L1010 ATCC 23745	64	> 128	> 128	> 128
<u>Haemophilus influenzae</u> type B	8	> 128	> 128	> 128
<u>Esch. coli</u> L47 SKF 12140	> 128	> 128	> 128	> 128
<u>Prot. vulgaris</u> ATCC 881	> 128	> 128	> 128	> 128

TABLE VII - (MIC, microgram/ml) (continued)

STRAIN	COMPOUND NO.					
	36	37	38	39	40	
<i>Staph. aureus</i> L165 Tour	0.13	0.03	32	0.13	0.25	
<i>Staph. epidermidis</i> L147 ATCC 12228	0.13	0.06	32	0.25	0.25	
<i>Staph. haemolyticus</i> L602	0.13	0.13	64	2	1	
<i>Strep. pneumoniae</i> L44 UC41	4	>128	>128	4	1	
<i>Strep. faecalis</i> L149 ATCC 7080	0.13	0.06	16	<0.13	<0.13	
<i>Prop. acnes</i> L1014 ATCC 6919	0.06	0.004	0.25	<0.13	.	
<i>Bact. fragilis</i> L1010 ATCC 23745	>128	>128	>128	8	4	
<i>Haemophilus influenzae</i> type B	>128	>128	>128	1	1	
<i>Esch. coli</i> L47 SKF 12140	>128	>128	>128	>128	>128	
<i>Prot. vulgaris</i> ATCC 881	>128	>128	>128	>128	>128	

TABLE VII - (MIC, microgram/ml) (continued)

STRAIN	COMPOUND NO.			
	41	42	43	44
<u>Staph. aureus</u> L165 Tour	0.25	0.13	0.13	0.13
<u>Staph. epidermidis</u> L147 ATCC 12228	0.5	0.13	0.5	0.5
<u>Staph. haemolyticus</u> L602	1	0.5	0.5	1
<u>Strep. pneumoniae</u> L44 UC41	2	1	0.5	1
<u>Strep. faecalis</u> L149 ATCC 7080	<0.13	0.13	0.13	0.06
<u>Prop. acnes</u> L1014 ATCC 6919	<0.13	0.016	0.016	0.016
<u>Bact. fragilis</u> L1010 ATCC 23745	4	>128	>128	>128
<u>Haemophilus Influenzae</u> type B	1	4	>128	>128
<u>Esch. coli</u> L47 SKF 12140	>128	>128	>128	>128
<u>Prot. vulgaris</u> ATCC 881	>128	>128	>128	>128

In view of their properties, the compounds of the invention can be used as active ingredients in the preparation of medicaments for human or animal treatment.

In particular, the amide derivatives of the antibiotic GE 2270 compounds of formula I are antimicrobial agents mainly active against gram positive bacteria and gram positive as well as gram negative anaerobes.

The main therapeutic indication of the antibiotic substances of the invention is thus in the treatment of infections related to the presence of microorganisms susceptible to them.

The term "treatment" is intended to encompass also prophylaxis, therapy and cure.

The patient receiving this treatment is any animal in need, including primates, in particular humans, and other mammals such as equines, cattle, swine and sheep; and poultry and pets in general.

The compounds of the invention can be administered as such or in admixture with pharmaceutically acceptable carriers and can also be administered in conjunction with other antimicrobial agents. Conjunctive therapy, thus includes sequential, simultaneous and separate administration of the active compounds in a way that the therapeutical effect of the first administered one is not entirely disappeared when the subsequent is administered.

A preferred pharmaceutical formulation is represented by a formulation suitable for a topical application on an intact or damaged skin or mucous membrane. Examples of such formulations are powders, ointments, creams and lotions. The excipients in these formulations are the usual pharmaceutically acceptable vehicles such oleaginous ointment bases (e.g. cetyl esters wax, oleic acid, olive oil, paraffin, spermaceti, starch glycerite); absorbent ointment bases (e.g. anhydrous lanolin, hydrophilic petrolatum), emulsion ointment bases (e.g. cetyl alcohol, glyceryl monostearate, lanolin, stearic acid), water-soluble ointment bases (e.g. glycol ethers and their derivatives which include polyethylene glycols, poly(oxy-1,2-ethanediyl)-alpha-hydro-omega-hydroxy-octadecanoate, polysorbates, and polyethylene glycols mono-stearates).

These formulations may contain other known excipients, such as preservatives and are prepared as known in the art and reported in reference handbooks such as Remington's Pharmaceutical Sciences, Seventeenth edition, 1985, Mack Publishing Co.

The compounds of the invention can also be formulated into formulation suitable for parenteral administration according to procedures known per se in the art. For instance, a compound of the invention is formulated with polypropylene glycol or dimethylacetamide and a surface-active agent such as polyoxyethylene sorbitan mono-oleate or polyethoxylated castor oil.

A preferred formulation for parenteral administration includes the following excipients: Cremophor® EL (polyoxyl 35 castor oil USP/NF) 20%, propylene glycol 5-10%.

Preferably, this formulation is used for i.v. administration in the treatment of any infection involving a microorganism susceptible to an antibiotic of the invention.

An example of a suitable formulation used for i.v. is the following

compound No. 19	100 mg
propylene glycol	1 ml
water for injection q.s.	5 ml
phosphate buffer pH 8-8.5	

In the treatment of pseudomembranous colitis or other diseases attributable to the presence of anaerobes in the gastrointestinal tract, an effective dose of the compounds of the invention may be administered orally in suitable pharmaceutical form such as a capsule or an aqueous suspension.

The dosage of the active ingredient depends on many factors which include type, age and conditions of the patient, specific active ingredient and formulation selected for administration, administration schedule, etc.

In general, effective antimicrobial dosages are employed per single unit dosage form. Repeated applications of these dosage forms, e.g. from 2 to 6 times a day, are in general preferred. An effective dosage may be in general in the range 0.5-50 mg/kg body weight/day.

A preferred topic preparation is an ointment containing from 1% to 10% of a compound of the present invention.

Anyway, the prescribing physician will be able to determine the optimal dosage for a given patient in a given situation.

Besides their use as medicaments in human and veterinary therapy, the compounds of the invention can also be used as animal growth promoters.

For this purpose, a compound of the invention is administered orally in a suitable feed. The exact concentration employed is that which is required to provide for the active agent in a growth promotant effective amount when normal amounts of feed are consumed.

The addition of the active compound of the invention to animal feed is preferably accomplished by preparing an appropriate feed premix containing the active compound in an effective amount and

incorporating the premix into the complete ration.

Alternatively, an intermediate concentrate or feed supplement containing the active ingredient can be blended into the feed. The way in which such feed premixes and complete rations can be prepared and administered is described in reference books (such as "Applied Animal Nutrition", W.H. Freedman and Co., S. Francisco, USA, 1969 or "Livestock Feeds and Feeding", O and B books, Corvallis, Oregon, USA, 1977).

The following examples further illustrate the invention and should not be interpreted as limiting it in any way.

EXAMPLES OF THE INVENTION

PROCEDURE A - Reaction of GE 2270 factor A₃ starting material with the selected amine

Example 1:

Preparation of compounds no. 15, 29, 30, 32, 33

To a stirred solution of 1 mmol of GE 2270 factor A₃ (prepared as described in European Patent Application Publication No. 406745) in 10 ml of dimethylformamide (DMF), 1.2 mmols of the selected amine, 1.4 mmols of triethylamine (TEA) and 1.2 mmols of diphenylphosphorazidate (DPPA) were added at 0 °C. (If the salt (chloride, p-toluenesulfonate, etc) of the selected amine was used, a double amount of TEA had to be used). The temperature was allowed to rise to room temperature and stirring was continued for about 4 h. The reaction mixture was then acidified with 1N aq HCl to about pH 3 and then diluted with water to complete precipitation of the product. The wet solid was dried in air and then purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 3 to 5% methanol in chloroform. Fractions containing the title compound were pooled together and the solvent evaporated. Trituration of the solid with ethyl ether yielded the title compound as a fine powder.

PROCEDURE A1 - Reaction of GE 2270 factor A₃ starting material with the selected amine containing further reactive functional group(s), all of which protected, and subsequent removal of the protecting group(s).

Example 2:

Preparation of compounds no. 34, 36

The reaction was substantially carried out as described in Example 1. Once the reaction product had been purified by flash chromatography, 1 mmol of the solid obtained was treated with 7 ml of cold trifluoroacetic acid (TFA). The suspension was swirled for a few minutes until a solution was obtained and TFA was evaporated "in vacuo" in the cold. The gummy product still containing traces of TFA was then treated with ethyl ether and the title compound was obtained as the trifluoroacetate salt in the form of a fine powder.

Example 3:

Preparation of compounds no. 1, 3 to 10, 18 to 21, 39

The reaction was substantially carried out as described in Example 1. Once the reaction product had been purified by flash chromatography, 1 mmol of the solid obtained was dissolved in 20 ml of dioxane and 1.2 ml of 1N aq NaOH were added under stirring at room temperature. After 5 h the solution was acidified with 1N aq HCl to pH 2 and diluted with water to complete precipitation of the title compound which was filtered off and allowed to dry in air yielding the title compound as a fine powder.

Example 4:

Preparation of compound no. 2

The reaction was carried out as described in Example 3. Once hydrolysis of the ester function had been accomplished and the compound had been allowed to dry in air, 1 mmols of the solid obtained was dissolved in 20 ml of TFA and 50 mmols of thioanisole were added under stirring at room temperature as

described by Y. Kiso et al., Chem. Pharm. Bull. 28, 673, 1980. After 3.5 h, TFA was evaporated "in vacuo" in the cold and the residue taken up in a minimum amount of 1% methanol in chloroform. Addition of ethyl ether induced the precipitation of the title compound which was filtered, washed with more ethyl ether and dried "in vacuo" to yield the trifluoroacetate salt of the title compound as a fine powder.

Example 4bis:

Preparation of compound no. 37

The reaction was substantially carried out as described in Example 1. Once the starting material had disappeared from the reaction mixture, water was added and the precipitate obtained was filtered off, washed with additional water and allowed to dry in air. The crude material was then dissolved in 3 ml of THF and stirred overnight at room temperature in the presence of 10% aq. HCl. Dilution with water provided complete precipitation of the product which was filtered off and allowed to dry in air. The solid was then purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 2 to 4% methanol in chloroform. Fractions containing the title compound were pooled together and the solvent evaporated yielding a pale yellow powder.

PROCEDURE B - Reaction of GE 2270 factor A₃ starting material with the selected amine containing unprotected acid moieties.

Example 5

Preparation of compounds no. 19, 22 to 28, 40, 41

1.1 mmols of DPPA were added at 0°C to a stirred solution of 1 mmol of GE 2270 factor A₃ and 1.5 mmols of TEA in 10 ml of DMF. The temperature was allowed to rise to room temperature and stirring was continued for 4.5 more hours. 1.5 mmols of the selected amino and 2 mmols of TEA were then added to the solution at room temperature and stirring was continued at the same temperature for 5 more hours. (If the selected amine contained more than one acid function, the amount of TEA was adjusted so to free the amino group). The reaction mixture was then acidified with 1N aq HCl to about pH 2 and then diluted with water to complete precipitation of the product. The wet solid was dried in air and then purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 5 to 10% methanol in chloroform. Fractions containing the title compound were pooled together and the solvent evaporated. Trituration of the solid with ethyl ether yielded the title compound as a fine powder.

PROCEDURE BI - Reaction of GE 2270 factor A₃ starting material with the selected amine containing reactive functional group(s), all of which are variously protected, in addition to the unprotected acid group(s) and subsequent removal of the protecting group(s).

Example 6:

Preparation of compounds no. 11, 12

The reaction was substantially carried out as described in Example 5. Once the reaction product had been purified by flash chromatography, 1 mmol of the solid obtained was dissolved in 20 ml of TFA and 50 mmols of thioanisole were added under stirring at room temperature. After 3.5 h, TFA was evaporated "in vacuo" in the cold and the residue taken up in a minimum-amount of 1% methanol in chloroform. Addition of ethyl ether induced the precipitation of the title compound which was filtered, washed with more ethyl ether and dried "in vacuo" to yield the trifluoroacetate salt of the title compound as a fine powder.

PROCEDURE C - Reaction of selected amide derivatives of GE 2270 factor A₃ as starting material with the selected reagent.

Example 7

Preparation of compounds no. 14, 15, 16, 17 from compounds no. 1, 5, 10, 6 respectively

To a stirred solution of 1 mmol of the appropriate amide derivative of GE 2270 factor A₃ (prepared as described in the previous examples) in 10 ml of DMF, 1.2 mmols of the selected amine, 1.4 mmols of TEA and 1.2 mmols of DPPA were added at 0°C. (If the salt (chloride, p-toluenesulfonate, etc) of the selected

amine was used, a double amount of TEA had to be used). The temperature was allowed to rise to room temperature and stirring was continued for about 4 h. The reaction mixture was then acidified with 1N aq HCl to about pH 3 and then diluted with water to complete precipitation of the product. The wet solid was dried in air and then purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 3 to 5% methanol in chloroform. Fractions containing the title compound were pooled together and the solvent evaporated. Trituration of the solid with ethyl ether yielded the title compound as a fine powder.

PROCEDURE CI - Reaction of the selected amide derivative of the GE 2270 factor A₃ as starting material with the selected reagent which contains further reactive functional group(s), all of which protected, and subsequent removal of the protecting group(s).

Example 8:

Preparation of compound no. 13 from compound no.3

The reaction was carried out as described in Example 7. Once the reaction product had been purified by flash chromatography, 1 mmol of the solid obtained was dissolved in 20 ml of dioxane and 1.2 ml of 1N aq NaOH were added under stirring at room temperature. After 5 h the solution was acidified with 1N aq HCl to pH 2 and diluted with water to complete precipitation of the title compound which was filtered off and allowed to dry in air yielding the title compound as a fine powder.

Example 9:

Preparation of compound no. 31 from compound no. 36

To a stirred solution of 1 mmol of the appropriate amide derivative of GE 2270 factor A₃ (prepared as described in the previous examples) in 10 ml of 10% methanolic chloroform, 1.2 mmols of TEA and 1.1 mmols of the selected reagent (see table II) were added at room temperature. After 20 min the solvent was evaporated "in vacuo" and the residue treated with 5% aq Na₂CO₃. The solid obtained was filtered off, washed with more 5% Na₂CO₃ and water and finally redissolved in 10 ml of methanol. To this solution, 0.5 ml of water and 0.1 mmols of p-toluenesulfonic acid were added and the reaction mixture was stirred at room temperature overnight. The solution was then reduced to a small volume (about 2 ml) under vacuum and water was added to precipitate the title compound which, after drying in air, was obtained as a fine powder.

Example 9bis:

Preparation of compound no. 38 from compound no. 37

To a stirred solution of 0.23 mmols of the appropriate amide derivative of GE 2270 factor A₃ (prepared as described in the previous examples) in 40 ml of ethanol, 9.2 mmols of acetic acid, 9.2 mmols of sodium acetate and 0.506 mmols of the selected reagent (see table II) were added at room temperature. After 2 hours 0.46 mmols of NaBH₄ (Fluka) were added and stirring was continued overnight at the same temperature. Evaporation of the solvent provided a crude material which was washed with 10 ml of 1N HCl, filtered and allowed to dry in air. The solid was then purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 0 to 10% methanol in dichloromethane. The fractions containing the methyl ester of the title compound (intermediate) were pooled together and the solvent evaporated providing a solid which was redissolved in 2 ml dioxane and treated overnight with a 1.2 molar excess of 1N NaOH at room temperature. Evaporation of the solvent gave a solid which was further purified by trituration with a 1:1 (v/v) mixture of ethyl acetate:methanol yielding the title compound as a fine powder.

PROCEDURE D - Reaction of GE 2270 factor A₂ starting material with the selected amine

Example 10:

Preparation of compound No. 35

1 mmol of GE 2270 factor A₂ (prepared as described in European Patent Application Publication No. 406745) was dissolved in 10 ml of a saturated solution of methanolic ammonia. The solution was allowed to

stand for 3 days at room temperature and then evaporated "in vacuo". The residue was taken up in 2 ml of methanol and the title compound precipitated with water, filtered off and allowed to dry in air. Trituration with ethyl ether yielded the title compound as a fine powder.

PROCEDURE E - Preparation of a salt of a compound of the invention.

5

Example 11:

Preparation of the arginine salt of compound No. 19

10 To a suspension of 3 g of compound No. 19 (2.42 mmols) in 180 ml of dioxane, a solution of 423 mg of L-arginine (2.42 mmols) in 120 ml of water were added under stirring and the non clear solution was thus lyophilized to recovered the desired salt.

PROCEDURE F - Reaction of GE 2270 factor C_{2a} starting material (i.e. the compound of formula II wherein R is methoxymethyl, R₁ is methyl, R₄ is hydroxymethyl and W is COOH) with the selected amine
15 containing further reactive functional group(s), all of which protected, and subsequent removal of the protecting group(s).

Example 12:

20 Preparation of compound no. 42

The reaction was carried out as described in Example 3 using GE 2270 factor C_{2a} starting material instead of factor A₃.

PROCEDURE G - Reaction of GE 2270 factor C_{2a} starting material as described in procedure F with the
25 selected amine containing unprotected acid moieties.

Example 13:

Preparation of compound no.42

30

The reaction was carried out as described in Example 5 using GE 2270 factor C_{2a} starting material instead of factor A₃.

PROCEDURE H - Reaction of GE 2270 factor D₁ starting material (i.e. the compound of formula II wherein R and R₁ are hydrogen, R₄ is methyl and W is COOH) with the selected amine containing further
35 reactive functional group(s), all of which protected, and subsequent removal of the protecting group(s).

Example 14:

Preparation of compound no. 43

40

The reaction was carried out as described in Example 3 using GE 2270 factor D₁ starting material instead of factor A₃.

PROCEDURE I - Reaction of GE 2270 factor D₁ starting material as described in procedure H with the
45 selected amine containing unprotected acid moieties.

Example 15:

Preparation of compound no.43

50

The reaction was carried out as described in Example 5 using GE 2270 factor D₁ starting material instead of factor A₃.

PROCEDURE J - Reaction of GE 2270 factor D₂ (i.e. the compound of formula II wherein R is hydroxymethyl, R₁ and R₄ are methyl and W is COOH) starting material with the selected amine containing
55 further reactive functional group(s), all of which protected, and subsequent removal of the protecting group(s).

Example 16:

Preparation of compound no. 44

- 5 The reaction was carried out as described in Example 3 using GE 2270 factor D₂ starting material instead of factor A₃.

PROCEDURE K - Reaction of GE 2270 factor D₂ starting material as described in procedure J with the selected amine containing unprotected acid moieties.

10 Example 17:

Preparation of compound no.44

- 15 The reaction was carried out as described in Example 5 using GE 2270 factor D₂ starting material instead of factor A₃.

PROCEDURE L - Reaction of a mixture of minor factors (C_{2a}, D₁, D₂ and E) of antibiotic GE 2270 (starting material) with the selected amine containing further reactive functional group(s), all of which protected, and subsequent removal of the protecting group(s).

20 Example 18:

The reaction was carried out as described in Example 3 using a mixture of minor components (C_{2a}, D₁, D₂ and E) of antibiotic GE 2270 starting material instead of factor A₃ and methyl 6-aminocaproate hydrochloride (Fluka). R_t (min) refer to HPLC method M reported in the HPLC analysis section.

- 25 When Y = -NHCH₂CH₂CH₂CH₂CH₂COOCH₃, R_t (min) are respectively 43.43 for GE 2270 factor C_{2a}, 39.42 for GE 2270 factor D₁, 42.29 for GE 2270 factor D₂ and 37.41 for GE 2270 factor E.

When Y = -NHCH₂CH₂CH₂CH₂CH₂COOH, R_t (min) are respectively 17.23 for GE 2270 factor C_{2a}, 15.76 for GE 2270 factor D₁, 16.64 for GE 2270 factor D₂ and 15.13 for GE 2270 factor E.

- 30 PROCEDURE M - Reaction of a selected mixture of minor components (C_{2a}, D₁, D₂ and E) of antibiotic GE 2270 starting material with the selected amine containing unprotected acid moieties.

Example 19:

- 35 The reaction was carried out as described in Example 5 using a selected mixture of minor components (C_{2a}, D₁, D₂ and E) of antibiotic GE 2270 starting material instead of factor A₃ and 6-aminocaproic acid (Fluka).

R_t (min) refer to Method M reported in the HPLC analysis section and are respectively 17.23 for GE 2270 factor C_{2a}, 15.76 for GE 2270 factor D₁, 16.64 for GE 2270 factor D₂ and 15.13 for GE 2270 factor E.

40 PREPARATION OF THE STARTING MATERIALS

1. The following starting materials have been purchased from Fluka (Fluka, Chemika-Biochemika, Buchs, Switzerland):

- 45 Glycine ethyl ester hydrochloride,
L-threonine methyl ester hydrochloride,
L-tyrosine methyl ester hydrochloride,
L-leucine methyl ester hydrochloride,
L-phenylalanine methyl ester hydrochloride,
L-methionine methyl ester hydrochloride,
50 L-proline methyl ester hydrochloride,
N α -Cbz-L-lysine,
methyl 4-aminobutyrate hydrochloride,
methyl 6-aminocaproate hydrochloride,
6-aminocaproic acid,
55 4-(methylamino)benzoic acid,
piperidine-4-carboxylic acid,
N-methyl-D-glucamine,
D(+) -glucosamine hydrochloride,

2-dimethylaminoethylamine,
amino acetaldehyde dimethylacetal,
 β -alanine ethyl ester hydrochloride.

2. The following starting materials have been purchased from Sigma (Sigma, Biochemicals Organic Compounds, St. Louis, U.S.A.):

N δ -Cbz-L-ornithine,
L-aspartic acid dimethyl ester hydrochloride.

3. The following starting materials have been purchased from Aldrich (Aldrich, Catalogo Prodotti di Chimica Fine, Milano, Italy):

L-Prolinamide,
taurine,
3-amino-1-propanesulfonic acid,
3-aminopropylphosphonic acid,
4-amino-1-benzylpiperidine.

4. Production of antibiotic GE 2270 for preparing antibiotic GE 2270 factors A, B₁, B₂, C₁, C₂, C_{2a}, D₁, D₂, and E

A culture of *Planobispora rosea* ATCC 53773 is grown on an oatmeal agar slant for two weeks at 28-30 °C and then used to inoculate 500 ml flasks containing 100 ml of a seed medium of the following composition:

Starch	20 g/l
Polypeptone	5 g/l
Yeast extract	3 g/l
Beef extract	2 g/l
Soybean meal	2 g/l
Calcium carbonate	1 g/l
Distilled water q.s.	100 ml
(adjusted to pH 7.0 before sterilization)	

The flask is incubated on a rotary shaker (200 rpm) at 28-30 °C for 92 h. The obtained culture is then used to inoculate a jar fermenter containing 4 liters of the same medium and the culture is incubated at 28-30 °C for 48 hours with stirring (about 900 rpm) and aeration (about one standard liter of air per volume per minute).

The obtained broth is transferred to a fermenter containing 50 l of the following production medium:

Starch	20 g/l
Peptone	2.5 g/l
Hydrolyzed casein	2.5 g/l
Yeast extract	3 g/l
Beef extract	2 g/l
Soybean meal	2 g/l
Calcium carbonate	1 g/l
Distilled water	q.s.
(adjusted to pH 7.0 before sterilization)	
and incubated for about 72 hours at 28-30 °C.	

Antibiotic production is monitored by paper disc agar assay using *B. subtilis* ATCC 6633 grown on minimum Davis medium. The inhibition zones are evaluated after incubation overnight at 35 °C.

4a) Recovery of crude antibiotic GE 2270

The fermentation mass (50 l) obtained above is harvested and submitted to filtration in the presence of a filter aid (Clarcell).

Antibiotic GE 2270 is found mainly in the mycelium, even if a certain amount of it can be recovered also from the filtrates.

The filtrate is adjusted to about pH 7.0 and extracted with ethyl acetate (50 l). The organic phase is separated by centrifugation and concentrated to a small volume under reduced pressure. The obtained oily residue is then treated with petroleum ether to precipitate crude antibiotic GE 2270 that is collected by filtration and dried. 415 mg of crude antibiotic GE 2270 complex is obtained.

The mycelium is extracted twice with 20 l of methanol and the pooled extracts are concentrated under reduced pressure to give an aqueous residue which is extracted twice with ethyl acetate. Crude antibiotic GE 2270 (6.06 g) is precipitated by addition of petroleum ether from the concentrated organic phase.

10

4b) Isolation of antibiotic GE 2270 factor A

The crude obtained from the mycelium according to the procedure described above (3 g) is dissolved in tetrahydrofuran and concentrated under reduced pressure in the presence of silica gel (230-400 mesh). The obtained solid residue is collected and applied to a chromatography column containing 300 g of silica gel (230-400 mesh) prepared in methylene chloride (CH_2Cl_2). The column is developed first with methylene chloride (2 l) and then sequentially with 1.5 l mixtures of methylene chloride and methanol in the following ratios: 98/2; 96/4, 94/6, 92/8, 90/10 and 88/12 (v/v).

Fractions are collected, analyzed by TLC, HPLC or microbiologically against *B. subtilis* and pooled according to their antibiotic content.

The pooled fractions containing antibiotic GE 2270 factor A are concentrated under reduced pressure to give an oily residue which is solubilized with tetrahydrofuran.

From this solution, antibiotic GE 2270 factor A (600 mg) is precipitated by adding petroleum ether.

25 4bis) Isolation of mixtures of minor components of antibiotic GE 2270

A representative mixture particularly enriched in the minor factors C_{2a} , D_1 , D_2 and E was established by HPLC comparison with analytical samples of each single component.

R_t (min) refer to HPLC method M reported in the HPLC analysis section and are 20.55 for GE 2270 factor C_{2a} , 17.43 for GE 2270 factor D_1 , 18.17 for GE 2270 factor D_2 , and 16.61 for GE 2270 factor E.

Concentration of this fraction under reduced pressure produced an oily residue which was redissolved in tetrahydrofuran and precipitated with petroleum ether as whitish powder.

35 4c) Separation and isolation of antibiotic GE 2270 factors D_1 , D_2 , and E

Antibiotic GE 2270 factors D_1 , D_2 and E are separated and purified from the above obtained crude mixture by preparative HPLC using a 250x20 mm column packed with Nucleosil® C18 (silica gel functionalized with octadecylsilane groups) (5 μm) and eluted with mixtures of Phase A: CH_3CN :tetrahydrofuran:40 mM HCOONH_4 (40:40:20) (v/v/v); Phase B: CH_3CN :tetrahydrofuran:40 mM HCOONH_4 (10:10:80) (v/v/v). The antibiotic mixture (6 mg) was solubilized in 3 ml of Phase B and 1 ml of Phase A and was injected into the HPLC column which was eluted at a flow rate of 14 ml/min with a 26:74 (v/v) mixture of Phase A and B. The eluted fractions were collected according to the UV absorption profile at 254 nm. The fractions of subsequent chromatographic runs having homogeneous content were pooled and concentrated under reduced pressure to eliminate CH_3CN . The residual solution showed antibacterial activity against *Staphylococcus aureus* Tour L165 by paper disc assay. These solutions were lyophilized at least three times to remove completely the HCOONH_4 buffer residue from the HPLC phases.

The yields were as follows: antibiotic GE 2270 factor E, 11 mg; antibiotic GE 2270 factor D_1 , 12 mg; antibiotic GE 2270 factor D_2 , 10 mg.

50 4d) Isolation of a purified mixture containing antibiotic GE 2270 factor C_{2a} and other GE 2270 factors

The preparations of crude GE 2270 factors from 6 repeated fermentations were pooled and solubilized into 12 l of CH_2Cl_2 :methanol (93:7) (v/v). The insoluble material was removed by filtration and the solution, containing the antibiotic complex, was applied to a 13 kg (230-400 mesh) silica gel column equilibrated in CH_2Cl_2 :methanol (93:7) (v/v). Antibiotic GE 2270 factor C_{2a} was eluted from the column by eluting with CH_2Cl_2 :methanol (93:7) (v/v). The fractions containing it (HPLC analysis) were pooled, were concentrated under reduced pressure and were dried to yield 23.5 g of antibiotic GE 2270 factor C_{2a} in mixture with other minor factors.

A portion (5.5 g) of this preparation was again purified by flash chromatography on a column containing 400 g of silica gel (230-400 mesh) equilibrated in methylene chloride (CH_2Cl_2). The column was developed first with methylene chloride (1 liter) and then sequentially with a series of mixtures of methylene chloride / methanol in the following ratios (v/v): 96/4 (3 liters); 94/6 (1 liter); 92/8 (2 liters); 90/10 (6 liters) and 88/12 (4 liters).

The fractions containing mainly GE 2270 factor C_{2a} (HPLC analysis) were pooled and were concentrated. The antibiotic preparation (646 mg) was precipitated upon addition of petroleum ether.

4e) Isolation of pure antibiotic GE 2270 factor C_{2a}

The purified mixture containing mainly antibiotic GE 2270 factor C_{2a} was further purified by preparative HPLC from the above described preparation.

A portion of the above described preparation of the antibiotic (10 mg) was solubilized in 1 ml of Phase A (CH_3CN : tetrahydrofuran : 40 mM HCOONH_4 - 40:40:20) (v/v/v) and 1 ml of Phase B (CH_3CN : tetrahydrofuran : 40 mM HCOONH_4 - 10:10:80) (v/v/v) and was injected into a HPLC 250x20 mm Hibar column (E. Merck; Darmstadt F.R. Germany) packed with 7 μm Nucleosil®C18 (silica gel functionalized with octadecylsilane groups) which was equilibrated with a mixture of 40% Phase A and 60% Phase B. The column was eluted at 15 ml/min flow rate with a 22 minutes linear gradient from 40% to 50% of Phase A. The UV detection was 254 nm. The fractions of 10 subsequent chromatographic runs containing the pure antibiotic of the title were pooled and were concentrated under reduced pressure to eliminate CH_3CN . Antibiotic GE 2270 factor C_{2a} precipitated from water. The precipitate was collected by centrifugation, was washed twice with distilled water and was dried under vacuum yielding 66 mg of the pure antibiotic.

5. Preparation of antibiotic GE 2270 factor A_2

Antibiotic GE 2270 factor A (prepared as described above) (86 mg) is dissolved in 17 ml of 95% ethanol and 1.7 ml of acetic acid. After incubation at 60 °C for 24 h, the resulting solution is diluted with 0.1M sodium phosphate buffer pH 7.5 (100 ml) and adjusted to pH 7.5 with 1M sodium hydroxide. Ethanol is removed by evaporation under reduced pressure and the aqueous residue is extracted twice with ethyl acetate (100 ml). The organic phase is concentrated under reduced pressure to obtain a solid residue which is solubilized with tetrahydrofuran and then precipitated by adding petroleum ether. Antibiotic GE 2270 factor A_2 (62 mg) is obtained with minor amounts of antibiotic GE 2270 factors A and A_1 . Pure antibiotic GE 2270 factor A_2 is obtained by preparative HPLC as follows:

10 mg of the above crude product is solubilized in tetrahydrofuran, diluted to the solubility limit with water and then injected into a HPLC system with a column (250 x 20 mm) packed with Nucleosil® C18 (5 micrometer) reverse phase silica gel by Stacroma®, eluting with a linear gradient from 64% to 93% of phase B in phase A, in 20 min, at a flow rate of about 15 ml/min. In this system, phase A is a 90:10 (v/v) mixture of 18 mM aqueous sodium phosphate pH 7.2 and acetonitrile, while phase B is a 40:60 (v/v) mixture of 18 mM aqueous sodium phosphate pH 7.2 and acetonitrile. Fractions of five consecutive runs are collected and UV monitored at 330 nm. Fractions which contain substantial amounts of antibiotic GE 2270 factor A_2 , which correspond to the major peaks of the UV-elution profile, are pooled and concentrated under reduced pressure to an aqueous phase which is extracted twice with ethyl acetate. This organic layer is then washed with distilled water to remove the residual inorganic salts and concentrated to precipitate a solid residue that is then dissolved in tetrahydrofuran and re-precipitated with petroleum ether, to obtain pure antibiotic GE 2270 factor A_2 (45 mg).

In European Patent Application Publication No. 406745 are described other alternative methods for preparing antibiotic GE 2270 factor A_2 from antibiotic GE 2270 factor A.

6. Preparation of antibiotic GE 2270 factor A_3

Antibiotic GE 2270 factor A_2 is incubated for 1 h at room temperature in 0.5M sodium carbonate. The reaction mixture is then diluted with cold water and brought to pH 6.5 with hydrochloric acid. The neutralized solution contains antibiotic GE 2270 factor A_3 as the main reaction product. This antibiotic is extracted from the aqueous phase with ethyl acetate and then is precipitated from the concentrated organic phase by adding petroleum ether.

Pure antibiotic GE 2270 factor A_3 is obtained by column chromatography as described below:

1.5 Grams of crude GE 2270 factor A_3 is dissolved in 60 ml of a 1/1 (v/v) mixture of methanol and dichloromethane and adsorbed on silica gel (75-230 mesh) by evaporation of the solvents under reduced

pressure. The solid residue is then put on the top of a silica gel (75-230 mesh) column (bed height 40 cm) equilibrated with dichloromethane. The column is then eluted with mixtures of methanol in dichloromethane in the order: 1) 2% methanol (450 ml); 2) 5% methanol (500 ml); 3) 10% methanol (600 ml); 4) 15% methanol (500 ml); 5) 20% methanol (500 ml); 6) 30% methanol (250 ml).

- 5 Fractions are collected and monitored by TLC and a microbiological assay on *B. subtilis* ATCC 6633. Antibiotic GE 2270 factor A₃ is normally present in the eluates which contain about 15-20% methanol.

The fractions containing the desired product are pooled and concentrated under reduced pressure. Upon addition of petroleum ether to the residue, antibiotic GE 2270 factor A₃ precipitates (854 mg of pure product).

10

7. Preparation of the proper starting material from antibiotic factors D₁, D₂, E and C_{2a}

- By substantially following the same procedure described at points 5 and 6 above but starting from the single factors D₁, D₂, E and C_{2a} of antibiotic GE 2270 instead of factor A, the proper starting materials of formula II wherein W is COOH or an activated ester, R is hydrogen, CH₂OCH₃ or CH₂OH, R₁ is CH₃ or hydrogen and R₄ is hydroxymethyl or methyl, are obtained.

7a) Preparation of proper starting material from a mixture of minor components (C_{2a}, D₁, D₂ and E) of antibiotic GE 2270

20

- By substantially following the same procedure described at point 5 and 6 above but starting from a mixture of minor components (C_{2a}, D₁, D₂ and E) of antibiotic GE 2270 instead of the single factor A, the proper starting material of formula II wherein W is COOH or an activated ester and R, R₁ and R₄ are respectively methoxymethyl, methyl and hydroxymethyl for C_{2a}, hydrogen, hydrogen and methyl for D₁, hydroxymethyl, methyl and methyl for D₂ and hydroxymethyl, hydrogen and methyl for E are obtained.

R₁ (min) refer to HPLC method M reported in the HPLC analysis section.

When W is an activated ester, R₁ (min) are respectively 22.51 for GE 2270 factor C_{2a}, 19.80 for GE 2270 factor D₁, 20.41 for GE 2270 factor D₂ and 18.92 for GE 2270 factor E.

- When W is COOH, R₁ (min) are respectively 12.99 for GE 2270 factor C_{2a}, 10.38 for GE 2270 factor D₁, 11.08 for GE 2270 factor D₂ and 9.03 for GE 2270 factor E.

8. Preparation of glycyL-Nε-Cbz-L-lysine trifluoroacetate

35

- 4.8 ml of DPPA (22 mmols) was added at 0°C to a well stirred solution of 3.5 g of BOC-glycine (Fluka) (20 mmols) and 7.28 g of Nε-Cbz-L-lysine methyl ester hydrochloride (Fluka) (22 mmols) in 50 ml of dry DMF. To this solution, a solution of 5.8 ml of TEA (42 mmols) in 50 ml of dry DMF was added at 0°C over a 10 - 15 min period. Stirring was continued for 2 more hours at 0°C and then overnight at room temperature. The reaction mixture was diluted with 250 ml of toluene and 500 ml of ethyl acetate and washed with 1N aq. HCl (x3), water, a saturated solution of NaHCO₃ and brine. Drying over Na₂SO₄ and evaporation of the solvent yielded 9.7 g of a thick oil which resisted any attempt of crystallization. NMR of this oil was in perfect agreement with the structure of BOC-glycyl-Nε-Cbz-L-lysine methyl ester.

- The oil was dissolved in 200 ml of acetone/dioxane 1:1 (v/v) and 22 ml of 1N aq. NaOH were added over a 30 min period at 0°C under stirring. The reaction mixture was then stirred for 45 min at room temperature, diluted with 300 ml of cold water, acidified with 25 ml of 1N aq HCl and extracted with chloroform (x3) and ethyl acetate (x3). Drying over Na₂SO₄ and evaporation of the solvent yielded 9.4 g of a gum which resisted any attempt of crystallization. NMR of this gum was in perfect agreement with the structure of BOC-glycyl-Nε-Cbz-L-lysine.

- The gummy compound was treated with 20 ml of cold trifluoroacetic acid (TFA). The reaction mixture was swirled at room temperature until all the compound went in solution. The solution was reduced to a small volume under vacuum in the cold and then ethyl ether was added to induce precipitation of the title compound. 9.6 g of glycyL-Nε-Cbz-L-lysine trifluoroacetate were obtained as a white powder. NMR was in perfect agreement with the structure.

55

9. Preparation of L-tyrosyl-L-prolinamide

0.48 ml of DPPA (2 mmols) were added at 0 °C to a well stirred solution of 538 mg of BOC-L-tyrosine (Fluka) (2 mmols), 228.3 mg of L-prolinamide (Aldrich) (2 mmols) and 168 mg of NaHCO₃ in 5 ml of dry DMF. The reaction mixture was stirred for 24 h at room temperature and then diluted with 50 ml of water and extracted with chloroform (x3). The organic phase was washed water, dried over Na₂SO₄ and the solvent evaporated to yield an oil which was purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with hexane/acetone 2:3 (v/v). 420 mg of BOC-L-tyrosyl-L-prolinamide were in this way obtained as a white solid. NMR was in agreement with the structure.

The solid obtained was dissolved in 6 ml of ethyl acetate and stirred for 48 h at room temperature in the presence of 4 ml of 3N aq. HCl. The reaction mixture was then evaporated to dryness in vacuo and the residue redissolved in ethanol was precipitated with ethyl ether. 302 mg of L-tyrosyl-L-prolinamide were obtained as a white powder. NMR was in perfect agreement with the structure.

10. Preparation of methyl 8-aminocaprylate and methyl 11-aminoundecanoate p-toluenesulfonates

A solution of 40 mmols of the selected amino acid (Fluka) and 15.2 g of p-toluenesulfonic acid monohydrate (Fluka) (80 mmols) in 200 ml of methanol was refluxed overnight. The solvent was then evaporated in vacuo and the residue redissolved in ethyl ether. After sometime the title compounds crystallized out quantitatively. The NMR of both compounds was in agreement with their structure.

11. Preparation of 5-aminopentylphosphonic acid

3.48 g of 5-amino-1-pentanol (Fluka) (33.7 mmols) and 5.0 g of phthalic anhydride (Fluka) (33.7 mmols) were melted together at 180 °C. This temperature was maintained for 90 min until no more water developed. The reaction mixture was allowed to cool to room temperature and the oily mixture was chromatographed on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 2% methanol in chloroform. 5.9 g of a pure oil were obtained. NMR was in agreement with the structure.

To the 5.9 g of the oily intermediate (25 mmols), 1.6 ml of PBr₃ (17 mmols) were added portionwise so to control the exothermic reaction. The reaction mixture was heated at 100 °C for 1.5 h and then poured into crushed ice. The solid material that separated was filtered and allowed to dry in air overnight. 6.6 g of the pure bromo intermediate were obtained. The mass was in agreement with the expected molecular weight.

500 mg of the pure bromo intermediate (1.69 mmols) and 140 mg of triethyl phosphite (Fluka) (0.84 mmols) were heated together at 150 °C for about 1 h. Other three portions of 140 mg of triethyl phosphite were then added at 30 min interval at the same temperature. When all the starting material had disappeared, the excess of triethyl phosphite was distilled off and the crude material purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 2% methanol in dichloromethane. 468 mg of the expected diethyl phosphonate were obtained as a thick oil. NMR confirmed the structure.

468 mg of the diethyl phosphonate intermediate were treated overnight with 3 ml of a 0.2 M solution of hydrazine in methanol at room temperature. The precipitated phthalhydrazide was filtered off and the remaining solution was evaporated to dryness in vacuo. The residue was taken up in 1 N aq. HCl and the solution was washed with ethyl acetate, basified with NaOH and extracted several times with n-butanol. The butanolic phase was dried over Na₂SO₄ and evaporated to dryness to yield 175 mg of a thick oil whose NMR was in agreement with the product expected.

175 mg of diethyl 5-aminopentylphosphonate were refluxed for 20 h in 0.6 ml of conc. HCl. The acid solution was then evaporated to dryness by azeotropic distillation in vacuo in the presence of n-butanol. The NMR of the glassy oil obtained confirmed it to be the 5-aminopentylphosphonic acid.

12. Preparation of 5-(5-aminopentyl)tetrazole

To a solution of 10 ml of 6-aminocapronitrile (Fluka) (80 mmols) and 13.3 ml of TEA (96 mmols) in 80 ml of tetrahydrofuran, 12.48 ml of benzyl chloroformate (Fluka) (88 mmols) were added dropwise at 0 °C under stirring. Stirring was continued for 2 h at room temperature and the solvent was evaporated in vacuo. The residue was dissolved in ethyl acetate, washed with 1 N aq. HCl, water and then dried over Na₂SO₄ and the solvent evaporated to yield 19.6 g of a syrup whose NMR was in agreement with the structure.

1 g of the protected 6-aminocapronitrile (4.06 mmols) in 40 ml of 1-methyl-2-pyrrolidone was heated at 150 °C under argon in the presence of 793 mg of sodium azide (12.2 mmols) and 834 mg triethylamine

hydrochloride (6.1 mmols). After 4 h the reaction mixture was diluted with 120 ml of water and then carefully acidified to pH 1 with 10% aq. HCl (attention: azotidric acid forms!). The solution was extracted with ethyl acetate, the organic phase re-extracted with 10% aq. NaOH (x2) and the basic solution washed with ethyl ether, acidified with conc. HCl and extracted with ethyl acetate (x3). Drying and evaporation of the organic phase yielded a syrup that crystallized from methanol/water. 260 mg of a fine powder were obtained. NMR and mass confirmed the structure.

250 mg of the N-protected aminotetrazole (0.86 mmols) were treated at room temperature with 5 ml of thioanisole (43.25 mmols) and 17.5 ml of trifluoroacetic acid for 3 h. Trifluoroacetic acid was concentrated in vacuo in the cold and ethyl ether was added to precipitate the title compound as its trifluoroacetate salt. NMR and mass confirmed the structure.

13. Preparation of N-[3,4-di-(O-tetrahydropyranyl)benzoyl]thiazolidin-2-thione

A solution of 4.62 g of 3,4-dihydroxybenzoic acid (Fluka) (30 mmols) in 40 ml of methanol was refluxed for 24 h in the presence of 0.325 ml of conc. H_2SO_4 . After cooling the solution to room temperature some solid NaHCO_3 was added and the solvent evaporated in vacuo. The residue was taken up in ethyl acetate, washed with water, dried over Na_2SO_4 and the solvent evaporated to yield a syrup which was crystallized from ethyl acetate/hexane. 3.53 g of white crystals were obtained.

9.1 ml of dihydropyran (Fluka) (0.1 mol) and 250 mg of pyridinium p-toluenesulfonate (1 mmol) were added at room temperature to a stirred solution of 1.68 g of methyl 3,4-dihydroxybenzoate (10 mmols) in 4 ml of ethyl acetate and 25 ml of dichloromethane. After 4 days the reaction mixture was washed with a saturated solution of NaHCO_3 , dried over Na_2SO_4 and evaporated to dryness to obtain 3.36 g of an oil which was used for the next step without further purification.

The crude from the previous reaction was dissolved in 40 ml of acetone and to the stirred solution 20 ml of water, 2.76 g of K_2CO_3 (20 mmols) and 10 ml of 1N aq. NaOH (10 mmols) were added and stirring was continued for 7 days at room temperature. Acetone was evaporated in vacuo and the residual water phase was washed with ethyl acetate. The aqueous phase was transferred to an flask containing an equal volume of chloroform, cooled to 0 °C and carefully acidified under vigorous stirring with 50 ml of 1 N aq. HCl. The water phase was then extracted 3 more times with chloroform and the combined organic layers were washed with 0.2% ammonium formate, dried over Na_2SO_4 and evaporated to dryness to yield a syrup which crystallized after hexane addition. 2.34 g of a white solid were obtained. The NMR was in agreement with the structure.

333 mg of 2-thiazoline-2-thiol (Fluka) (2.8 mmols), 577 mg of N,N'-dicyclohexylcarbodiimide (Fluka) (2.8 mmols) and 35 mg of 4-dimethylaminopyridine were added in the order at 0 °C to a stirred solution of 644 mg of the benzoic acid intermediate (2 mmols) in 14 ml ethyl acetate/dichloromethane 5:2 (v/v). Stirring was continued overnight at room temperature, the precipitated dicyclohexylurea was filtered off and the yellow solution was evaporated in vacuo to yield a yellow oil which was purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting from 25% acetone in hexane. 700 mg of yellow crystals were obtained from acetone/hexane. NMR and IR confirmed the compound to be the title compound.

14. Preparation of N¹,N⁸-di-tert-butoxycarbonylspermidine

A solution of 19.72 g of BOC-ON (Aldrich) (80 mmols) in 60 ml of degassed tetrahydrofuran (THF) was added dropwise over a 1 h period under argon to a stirred solution of 5.8 g of spermidine (Aldrich) (40 mmols) in 40 ml of degassed THF cooled at 0 °C. The reaction mixture was then stirred at room temperature overnight and then evaporated to dryness. The residue was taken up in ethyl ether, washed with 1 N aq. NaOH (x4) and water (x4), dried over Na_2SO_4 and the solvent concentrated to a small volume in vacuo. Upon addition of ethyl ether 11 g of a white powder precipitated. NMR confirmed it to be the title compound.

15. Preparation of N-tert-butoxycarbonylpropylenediamine

4.2 g of BOC-ON (Aldrich) (17.2 mmols) were added at room temperature to a stirred solution of 2 g of 3-aminopropionitrile fumarate (Aldrich) (15.6 mmols) dissolved in a mixture of 10 ml of dioxane, 10 ml of water and 3.3 ml of triethylamine. After 3 h the reaction mixture was diluted with more water and extracted with dichloromethane (x3). The combined organic layers were washed with 1 N aq. NaOH (x3) and water (x3), dried over Na_2SO_4 and evaporated to dryness. The residual oil was taken up in ethyl ether and precipitated with hexane to yield 2.2 g of a white powder.

1 g of N-BOC-protected intermediate (5.9 mmols) in 7 ml of 1 N ethanolic NaOH was hydrogenated at 40 psi in the presence of 130 mg of Raney nickel (50% slurry in water, pH>9) (Aldrich) for 40 h. Raney nickel was filtered off and the solvent was evaporated to dryness. The residue was taken up in ethyl acetate and washed with 1 N aq. NaOH, dried over Na₂SO₄ and the solvent removed in vacuo yielding 950 mg of a colorless oil which solidified on standing. NMR confirmed it to be the title compound.

16. Preparation of 3-(2-aminoethylthio)propanoic acid methyl ester trifluoroacetate

To a solution of 0.5 g of cysteamine (Fluka) (6.48 mmols) in 5 ml of CH₂Cl₂, 1.4 g of di-tert-butyl dicarbonate (Aldrich) (6.48 mmols) in 5 ml of CH₂Cl₂ were added at room temperature under stirring. After 30 min the organic solvent was evaporated and the crude material dissolved in 5 ml of absolute ethanol. To the ethanolic solution, 2.7 ml of TEA (19.1 mmols) and 1.07 ml of methyl 3-bromopropionate (Fluka) (9.57 mmols) were added in the order. The reaction was completed in about 30 min. Ethanol was removed in vacuo and replaced by 15 ml of chloroform. The organic phase was then washed with water, anhydriified on Na₂SO₄ and the solvent evaporated to yield an oil which was finally treated with 1 ml of trifluoroacetic acid at 0 °C for 5 min. Evaporation to dryness gave 270 mg of a pale yellow oil. NMR and IR confirmed it to be the title compound.

17. Preparation of 6-amino-2(E)-hexenoic acid

To a stirred solution of 2 ml of 4-amino-butyraldehyde diethyl acetal (Fluka) (11.6 mmols) and 3.6 ml of TEA (25.6 mmols) in 5 ml of CH₂Cl₂, a solution of 1.5 ml of benzoyl chloride (Fluka) (12.9 mmols) in 5 ml CH₂Cl₂ was added in 30 min. at room temperature. After 1 hour the reaction mixture was diluted with 10 more ml of CH₂Cl₂, washed with water and the organic phase dried over Na₂SO₄ and the volume adjusted to 20 ml. The new solution was allowed to react for three days under argon in the presence of 1.6 ml of TEA (11.5 mmols), 10.2 g of di-tert-butyl dicarbonate (Aldrich) (46.8 mmols) and 1.4 g of 4-dimethylaminopyridine (Fluka) (11.5 mmols) at room temperature. Removal of the solvent gave a brown oil that was purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 20% ethyl acetate in n-hexane yielding 1.6 g of the N,N deprotected 4-aminobutyraldehyde diethyl acetal as a colorless oil. NMR confirmed the structure.

The obtained oil was then dissolved in 5 ml of THF and treated with 5 ml of 1N HCl at room temperature for three hours. THF was removed in vacuo and the remaining solution was washed with chloroform (2 ml x3). The organic phase was then washed with a solution of Na₂CO₃, water, dried over Na₂SO₄ and evaporated to dryness yielding an oil that was used in the next step without further purification.

To a suspension of 160 mg of 60% NaH (4mmols) in 5 ml of dry THF at 0 °C under argon, 0.837 ml of triethylphosphonoacetate (Fluka) (4.3 mmols) were added. After 30 min. a dry THF (2 ml) solution of the previously obtained aldehyde (1.17 g) (4.02 mmols) was added and the temperature was allowed to rise to room temperature. The reaction mixture was stirred overnight and then 50 more mg of 60% NaH were added at 0 °C. After two more hours at room temperature the reaction mixture was treated with diluted HCl (10 ml) and extracted with ethyl acetate (5 ml x3). The combined organic phase was washed with water, dried over Na₂SO₄ and evaporated to dryness. The crude material was purified by flash chromatography on silica gel 60 (230 - 400 mesh ASTM - Merck) eluting with 15% ethyl acetate in n-hexane yielding 765 mg of a syrup. NMR confirmed it to be the expected product with the double bond in E configuration (J = 16Hz).

6.15 ml of 1N LiOH (6.15 mmols) were added to a solution of 739 mg of the unsaturated ester previously obtained (2.05 mmols) in 10 ml of THF under stirring at room temperature. When the starting material had disappeared the reaction mixture was concentrated in vacuo at 30 °C (bath temperature). The aqueous solution was acidified at pH 2 with 1N HCl and then extracted with ethyl acetate. The combined organic phase was dried over Na₂SO₄, filtered and the solvent evaporated yielding an oil that solidified upon standing under vacuum. NMR and MS confirmed it to be 6-N-BOC-amino-2(E)-hexenoic acid.

Removal of the N-BOC protection to obtain the title compound was carried out in neat trifluoroacetic acid at 0 °C just before the coupling with the appropriate GE 2270 starting material.

18. Preparation of 3-(2-aminoethoxy)propanoic acid trifluoroacetate


To a stirred solution of 1 g of N-BOC-ethanolamine (6.22 mmols) [prepared according to classical methodologies from ethanolamine (Fluka)] in 10 ml of dry THF at -78 °C, 3.88 ml of 1.6 M solution of butyllithium (Fluka) (6.22 mmols) were added under argon. After 30 min. 1.3 g of t-butyl 3-bromo propanoate [prepared according to classical methodologies from 3-bromo propanoic acid (Fluka)] (6.22

The total deprotection of the coupled compound was carried out immediately before addition to the appropriate GE 2270 starting material by stirring it in trifluoroacetic acid for about 5 min at room temperature. Removal of trifluoroacetic acid in vacuo yielded the title compound.

10 Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, MC, NL, SE, PT

1. An amide derivative of antibiotic GE 2270 having the following formula I




wherein

R represents:
hydrogen,
hydroxymethyl, or
methoxymethyl;

R₁ represents:
hydrogen, or
methyl:

Y represents:
a group of formula



5

10 wherein:

R₂

represents:

hydrogen,

(C₁-C₄)alkyl,amino(C₂-C₄)alkyl,

15

(C₁-C₄)alkylamino-(C₁-C₄)alkyl, ordi-(C₁-C₄)alkylamino-(C₁-C₄)alkyl;R₃

represents:

hydrogen,

20

a linear or branched (C₁-C₁₄)alkyl group bearing from 1 to 3 substituents selected from: carboxy, sulfo, phosphono, amino which may be optionally protected with a lower alkoxycarbonyl or a benzyloxycarbonyl group, (C₁-C₄)alkylamino wherein the alkyl moiety may be optionally substituted with a carboxy group,

di-(C₁-C₄)alkylamino, hydroxy, halo,

25

(C₁-C₄)alkoxy wherein the alkyl moiety may be optionally substituted with a carboxy group,

(C₁-C₄)alkoxycarbonyl, mercapto,

(C₁-C₄)alkylthio wherein the alkyl moiety may be optionally substituted with a carboxy group, phenyl which may be optionally substituted with 1 to 3 substituents selected from carboxy, sulfo, hydroxy, halo and mercapto, carbamyl,

30

(C₁-C₆)alkylcarbamyl wherein the alkyl moiety may be optionally substituted with 1 or 2 substituents selected from carboxy, amino,

(C₁-C₄)alkylamino and di-(C₁-C₄)alkylamino,

35

di-(C₁-C₄)alkylcarbamyl wherein the alkyl moieties together with the adjacent nitrogen atom may also represent a saturated 5-7 membered heterocyclic ring which may optionally be substituted with a carboxy or a carbamyl group on one of the ring carbons and may optionally contain a further heterogroup selected from O, S and N, benzoylamino wherein the phenyl group may be substituted from 1 to 3 hydroxy group, a nitrogen containing 5-6 membered heterocyclic ring which may be unsaturated, partially saturated or wholly saturated and may contain 1 to 3 further heteroatoms selected from N, S and O wherein one of the carbons of the ring may optionally bear a group carboxy, sulfo, carboxy(C₁-C₄)alkyl or sulfo(C₁-C₄)alkyl and the ring nitrogen atom may optionally be substituted by (C₁-C₄)alkyl, carboxy(C₁-C₄)alkyl, sulfo(C₁-C₄)alkyl, or benzyl;

40

(C₃-C₆)alkenyl, optionally substituted by carboxy or sulfo;

45

1-deoxy-1-glucityl;

2-deoxy-2-glucosyl;

a fully saturated 5 to 7 membered nitrogen containing heterocyclic ring wherein the nitrogen atom may be optionally substituted by (C₁-C₄)alkyl or benzyl and one or two carbons of the ring skeleton may bear a substituent selected from (C₁-C₄)alkyl, carboxy and sulfo;

50

or R₂ and R₃

taken together with the adjacent nitrogen atom represent a fully saturated 5-7 membered heterocyclic ring which may optionally contain a further heteroatom selected from O, S and N, and may optionally bear one or two substituents on the ring carbons selected from (C₁-C₄)alkyl, benzyl, carboxy, sulfo, carboxy(C₁-C₄)alkyl, and sulfo(C₁-C₄)alkyl;

55

R₄

represents:

hydrogen,

methyl, or

hydroxymethyl;

with the proviso that when R_4 is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and R_1 is methyl;

and the pharmaceutically acceptable addition salts thereof.


5

2. A compound according to claim 1 wherein R represents methoxymethyl and the other substituents are defined as in claim 1.

10

3. A compound as claimed in claim 1 wherein R represents methoxymethyl, R_1 and R_4 represent methyl and Y represents a group of formula

15



20

wherein R_2 is hydrogen and R_3 is defined as in claim 1.

4. A compound as claimed in claim 1 wherein R is methoxymethyl, R_1 and R_4 represent a methyl group and Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.

25


5. A compound as claimed in claim 1 wherein R is methoxymethyl, R_1 and R_4 are methyl, Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH , SO_3H and PO_3H_2 .

30

6. A compound as claimed in claim 1 wherein R is methoxymethyl, R_1 and R_4 are methyl, Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-COOH}$.

7. A compound as claimed in claim 1 wherein R represents hydrogen, hydroxymethyl or methoxymethyl, R_1 represents hydrogen or methyl, R_4 represents hydrogen, methyl or hydroxymethyl and Y represents a group of formula

40



45

wherein R_2 is hydrogen and R_3 is defined as in claim 1.

8. A compound as claimed in claim 7 wherein Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.


50

9. A compound as claimed in claim 1 wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 is hydrogen or methyl, R_4 is hydrogen, methyl or hydroxymethyl and Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH , SO_3H and PO_3H_2 .

55

10. A compound as claimed in claim 1 wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 is hydrogen or methyl, R_4 is hydrogen, methyl or hydroxymethyl and Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is $CH_2CH_2CH_2CH_2CH_2COOH$.

11. A process for preparing a compound of claim 1 which comprises reacting an antibiotic GE 2270 compound having formula II:



wherein

- 35 W represents a carboxy or an activated ester function;
 R represents hydrogen, hydroxymethyl or methoxymethyl;
 R_1 represents hydrogen or methyl;
 R_4 represents hydrogen, methyl or hydroxymethyl;

with the proviso that when R_4 is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and R_1 is methyl, with a selected amine of formula HNR_2R_3 wherein R_2 and R_3 have the same meanings as in claim 1, in an inert organic solvent and, when W is carboxy, in the presence of a condensing agent.

40

12. A process according to claim 11 wherein the condensing agents are selected from (C_1-C_4) alkyl, phenyl or heterocyclic phosphorazidates such as, diphenylphosphorazidate (DPPA), diethylphosphorazidate, di-(4-nitrophenyl)phosphorazidate, dimorpholylphosphorazidate and diphenylphosphorochloridate, or benzotriazol-1-yl-oxy-trispyrrolidinophosphoniumhexafluorophosphate (PyBOP).

45

13. A process according to claims 11 and 12 wherein the amine reactant HNR_2R_3 is used in a 1 to 2 fold molar excess with respect to the antibiotic starting material and the reaction temperature is comprised between 0 and 20 °C.

50

14. A compound of any of claims 1 to 10 for use as a medicine.

15. A pharmaceutical composition containing a compound of any of claims 1 to 10 as the active ingredient in admixture with a pharmaceutically acceptable carrier.

55

16. Use of a compound according to any of claims 1 to 10 for preparing a medicament for use as an antibiotic.

Claims for the following Contracting State : ES

1. A process for preparing an amide derivative of antibiotic GE 2270 having the following formula I

5

10

15

20

25

30

35

wherein

40 R represents:
hydrogen,
hydroxymethyl, or
methoxymethyl;

R₁ represents:
hydrogen, or
methyl;


45 Y represents:
a group of formula

50

55


wherein:

R₂ represents:



hydrogen,
 (C₁-C₄)alkyl,
 amino(C₂-C₄)alkyl,
 (C₁-C₄)alkylamino-(C₁-C₄)alkyl, or
 di-(C₁-C₄)alkylamino-(C₁-C₄)alkyl;
 5 R₃ represents:
 hydrogen,
 a linear or branched (C₁-C₁₄)alkyl group bearing from 1 to 3 substituents selected
 from: carboxy, sulfo, phosphono, amino which may be optionally protected with a
 10 lower alkoxy carbonyl or a benzyloxy carbonyl group, (C₁-C₄)alkylamino wherein the
 alkyl moiety may be optionally substituted with a carboxy group,
 di-(C₁-C₄)alkylamino, hydroxy, halo,
 (C₁-C₄)alkoxy wherein the alkyl moiety may be optionally substituted with a
 carboxy group,
 15 (C₁-C₄)alkoxy carbonyl, mercapto,
 (C₁-C₄)alkylthio wherein the alkyl moiety may be optionally substituted with a
 carboxy group, phenyl which may be optionally substituted with 1 to 3 substituents
 selected from carboxy, sulfo, hydroxy, halo and mercapto, carbamyl,
 (C₁-C₆)alkylcarbamyl wherein the alkyl moiety may be optionally substituted with 1
 20 or 2 substituents selected from carboxy, amino,
 (C₁-C₄)alkylamino and di-(C₁-C₄)alkylamino,
 di-(C₁-C₄)alkylcarbamyl wherein the alkyl moieties together with the adjacent
 nitrogen atom may also represent a saturated 5-7 membered heterocyclic ring
 which may optionally be substituted with a carboxy or a carbamyl group on one of
 25 the ring carbons and may optionally contain a further heterogroup selected from O,
 S and N, benzoylamino wherein the phenyl group may be substituted from 1 to 3
 hydroxy group, a nitrogen containing 5-6 membered heterocyclic ring which may
 be unsaturated, partially saturated or wholly saturated and may contain 1 to 3
 further heteroatoms selected from N, S and O wherein one of the carbons of the
 30 ring may optionally bear a group carboxy, sulfo, carboxy(C₁-C₄)alkyl or sulfo(C₁-
 C₄)alkyl and the ring nitrogen atom may optionally be substituted by (C₁-C₄)alkyl,
 carboxy(C₁-C₄)alkyl, sulfo(C₁-C₄)alkyl, or benzyl;
 (C₃-C₆)alkenyl, optionally substituted by carboxy or sulfo;
 1-deoxy-1-glucityl;
 35 2-deoxy-2-glucosyl;
 a fully saturated 5 to 7 membered nitrogen containing heterocyclic ring wherein the
 nitrogen atom may be optionally substituted by (C₁-C₄)alkyl or benzyl and one or
 two carbons of the ring skeleton may bear a substituent selected from (C₁-C₄)alkyl,
 carboxy and sulfo;
 40 or R₂ and R₃ taken together with the adjacent nitrogen atom represent a fully saturated 5-7
 membered heterocyclic ring which may optionally contain a further heteroatom
 selected from O, S and N, and may optionally bear one or two substituents on the
 ring carbons selected from (C₁-C₄)alkyl, benzyl, carboxy, sulfo, carboxy(C₁-C₄)
 alkyl, and sulfo(C₁-C₄)alkyl;
 45 R₄ represents:
 hydrogen,
 methyl, or
 hydroxymethyl;
 with the proviso that when R₄ is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl
 50 and R₁ is methyl;
 and the pharmaceutically acceptable addition salts thereof,
 which comprises reacting an antibiotic GE 2270 compound having formula II:

55



II

wherein

W represents a carboxy or an activated ester function;


R represents hydrogen, hydroxymethyl or methoxymethyl;

R₁ represents hydrogen or methyl;

R₄ represents hydrogen, methyl or hydroxymethyl;

with the proviso that when R₄ is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and R₁ is methyl, with a selected amine of formula HNR₂R₃ wherein R₂ and R₃ have the same meanings as above, in an inert organic solvent and, when W is carboxy, in the presence of a condensing agent.


2. A process according to claim 1 for preparing a compound of formula I wherein R represents methoxymethyl and the other substituents are defined as in claim 1.
3. A process according to claim 1 for preparing a compound of formula I wherein R represents methoxymethyl, R₁ and R₄ represent methyl and Y represents a group of formula



wherein R₂ is hydrogen and R₃ is defined as in claim 1.

4. A process according to claim 1 for preparing a compound of formula I wherein R is methoxymethyl, R₁ and R₄ represent a methyl group and Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.

5. A process according to claim 1 for preparing a compound of formula I wherein R is methoxymethyl, R₁ and R₄ are methyl, Y is a group NR₂R₃ wherein R₂ is hydrogen and R₃ is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH, SO₃H and PO₃H₂.
6. A process according to claim 1 for preparing a compound of formula I wherein R is methoxymethyl, R₁ and R₄ are methyl, Y is a group NR₂R₃ wherein R₂ is hydrogen and R₃ is CH₂CH₂CH₂CH₂CH₂-COOH.
7. A process according to claim 1 for preparing a compound of formula I wherein R represents hydrogen, hydroxymethyl or methoxymethyl, R₁ represents hydrogen or methyl, R₄ represents hydrogen, methyl or hydroxymethyl and Y represents a group of formula




wherein R₂ is hydrogen and R₃ is defined as in claim 1.

8. A process according to claim 1 for preparing a compound of formula I wherein Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.
9. A process according to claim 1 for preparing a compound of formula I wherein R is hydrogen, hydroxymethyl or methoxymethyl, R₁ is hydrogen or methyl, R₄ is hydrogen, methyl or hydroxymethyl and Y is a group NR₂R₃ wherein R₂ is hydrogen and R₃ is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH, SO₃H and PO₃H₂.
10. A process according to claim 1 for preparing a compound of formula I wherein R is hydrogen, hydroxymethyl or methoxymethyl, R₁ is hydrogen or methyl, R₄ is hydrogen, methyl or hydroxymethyl and Y is a group NR₂R₃ wherein R₂ is hydrogen and R₃ is CH₂CH₂CH₂CH₂CH₂-COOH.
11. A process according to anyone of claims 1 to 10 wherein the condensing agents are selected from (C₁-C₄)alkyl, phenyl or heterocyclic phosphorazidates such as, diphenylphosphorazidate (DPPA), diethylphosphorazidate, di(4-nitrophenyl)phosphorazidate, dimorpholylphosphorazidate and diphenylphosphorochloridate, or benzotriazol-1-yl-oxy-trisoyrolidinophosphoniumhexafluorophosphate (PyBOP).
12. A process according to anyone of claims 1 to 11 wherein the amine reactant HNR₂R₃ is used in a 1 to 2 fold molar excess with respect to the antibiotic starting material and the reaction temperature is comprised between 0 and 20 °C.
13. Use of a compound prepared according to any of claims 1 to 12 for preparing a medicament for use as an antibiotic.

- hydrogen,
 (C₁-C₄)alkyl,
 amino(C₂-C₄)alkyl,
 (C₁-C₄)alkylamino-(C₁-C₄)alkyl, or
 di-(C₁-C₄)alkylamino-(C₁-C₄)alkyl;
 5 R_3 represents:
 hydrogen,
 a linear or branched (C₁-C₁₄)alkyl group bearing from 1 to 3 substituents selected
 from: carboxy, sulfo, phosphono, amino which may be optionally protected with a
 10 lower alkoxy carbonyl or a benzyloxy carbonyl group, (C₁-C₄)alkylamino wherein
 the alkyl moiety may be optionally substituted with a carboxy group,
 di-(C₁-C₄)alkylamino, hydroxy, halo,
 (C₁-C₄)alkoxy wherein the alkyl moiety may be optionally substituted with a
 carboxy group,
 15 (C₁-C₄)alkoxycarbonyl, mercapto,
 (C₁-C₄)alkylthio wherein the alkyl moiety may be optionally substituted with a
 carboxy group, phenyl which may be optionally substituted with 1 to 3 substituents
 selected from carboxy, sulfo, hydroxy, halo and mercapto, carbamyl,
 (C₁-C₆)alkylcarbamyl wherein the alkyl moiety may be optionally substituted with 1
 20 or 2 substituents selected from carboxy, amino,
 (C₁-C₄)alkylamino and di-(C₁-C₄)alkylamino,
 di-(C₁-C₄)alkylcarbamyl wherein the alkyl moieties together with the adjacent
 nitrogen atom may also represent a saturated 5-7 membered heterocyclic ring
 which may optionally be substituted with a carboxy or a carbamyl group on one of
 25 the ring carbons and may optionally contain a further heterogroup selected from O,
 S and N, benzoylamino wherein the phenyl group may be substituted from 1 to 3
 hydroxy group, a nitrogen containing 5-6 membered heterocyclic ring which may
 be unsaturated, partially saturated or wholly saturated and may contain 1 to 3
 30 further heteroatoms selected from N, S and O wherein one of the carbons of the
 ring may optionally bear a group carboxy, sulfo,
 carboxy(C₁-C₄)alkyl or sulfo(C₁-C₄)alkyl and the ring nitrogen atom may optionally
 be substituted by (C₁-C₄)alkyl,
 carboxy(C₁-C₄)alkyl, sulfo(C₁-C₄)alkyl, or benzyl;
 (C₃-C₆)alkenyl, optionally substituted by carboxy or sulfo;
 35 1-deoxy-1-glucityl;
 2-deoxy-2-glucosyl;
 a fully saturated 5 to 7 membered nitrogen containing heterocyclic ring wherein the
 nitrogen atom may be optionally substituted by (C₁-C₄)alkyl of benzyl and one or
 two carbons of the ring skeleton may bear a substituent selected from (C₁-C₄)alkyl,
 40 carboxy and sulfo;
 R_2 and R_3 taken together with the adjacent nitrogen atom represent a fully saturated 5-7
 membered heterocyclic ring which may optionally contain a further heteroatom
 selected from O, S and N, and may optionally bear one of two substituents on the
 ring carbons selected from (C₁-C₄)alkyl, benzyl, carboxy, sulfo, carboxy(C₁-C₄)-
 45 alkyl, and sulfo(C₁-C₄)alkyl;
 R_4 represents:
 hydrogen,
 methyl, or
 hydroxymethyl;
 50 with the proviso that when R_4 is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl
 and R_1 is methyl;
 and the pharmaceutically acceptable addition salts thereof.

2. A compound according to claim 1 wherein R represents methoxymethyl and the other substituents are
 55 defined as in claim 1.
3. A compound as claimed in claim 1 wherein R represents methoxymethyl, R_1 and R_4 represent methyl
 and Y represents a group of formula




5

10 wherein R_2 is hydrogen and R_3 is defined as in claim 1.

4. A compound as claimed in claim 1 wherein R is methoxymethyl, R_1 and R_4 represent a methyl group and Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.
- 15
5. A compound as claimed in claim 1 wherein R is methoxymethyl, R_1 and R_4 are methyl, Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH , SO_3H and PO_3H_2 .
- 20
6. A compound as claimed in claim 1 wherein R is methoxymethyl, R_1 and R_4 are methyl, Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-COOH}$.
- 25
7. A compound as claimed in claim 1 wherein R represents hydrogen, hydroxymethyl or methoxymethyl, R_1 represents hydrogen or methyl, R_4 represents hydrogen, methyl or hydroxymethyl and Y represents a group of formula

30




35

wherein R_2 is hydrogen and R_3 is defined as in claim 1.

8. A compound as claimed in claim 7 wherein Y is an amino moiety which derives from a natural amino acid such as for example glycine, ornithine, serine, aspartic acid, tyrosine, leucine, phenylalanine, methionine, proline, threonine, lysine, or a synthetic dipeptide such as glycyllysine, serylproline, glycylprolinamide, tyrosylprolinamide, threonylprolinamide, leucylprolinamide.
- 40
9. A compound as claimed in claim 1 wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 is hydrogen or methyl, R_4 is hydrogen, methyl or hydroxymethyl and Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is a linear alkyl chain preferably of 3 to 12 carbons, more preferably of 3 to 7 carbons substituted with a group selected from COOH , SO_3H and PO_3H_2 .
- 45
10. A compound as claimed in claim 1 wherein R is hydrogen, hydroxymethyl or methoxymethyl, R_1 is hydrogen or methyl, R_4 is hydrogen, methyl or hydroxymethyl and Y is a group NR_2R_3 wherein R_2 is hydrogen and R_3 is $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-COOH}$.
- 50

55



93

R₃di-(C₁-C₄)alkylamino-(C₁-C₄)alkyl;

represents:

hydrogen,

a linear or branched (C₁-C₁₄)alkyl group bearing from 1 to 3 substituents selected from: carboxy, sulfo, phosphono, amino which may be optionally protected with a lower alkoxy carbonyl or a benzyloxy carbonyl group, (C₁-C₄)alkylamino wherein the alkyl moiety may be optionally substituted with a carboxy group,

di-(C₁-C₄)alkylamino, hydroxy, halo,

(C₁-C₄)alkoxy wherein the alkyl moiety may be optionally substituted with a carboxy group,

(C₁-C₄)alkoxycarbonyl, mercapto,

(C₁-C₄)alkylthio wherein the alkyl moiety may be optionally substituted with a carboxy group, phenyl which may be optionally substituted with 1 to 3 substituents selected from carboxy, sulfo, hydroxy, halo and mercapto, carbamyl,

(C₁-C₆)alkylcarbamyl wherein the alkyl moiety may be optionally substituted with 1 or 2 substituents selected from carboxy, amino,

(C₁-C₄)alkylamino and di-(C₁-C₄)alkylamino,

di-(C₁-C₄)alkylcarbamyl wherein the alkyl moieties together with the adjacent nitrogen atom may also represent a saturated 5-7 membered heterocyclic ring which may optionally be substituted with a carboxy or a carbamyl group on one of the ring carbons and may optionally contain a further heterogroup selected from O, S and N, benzoylamino wherein the phenyl group may be substituted from 1 to 3 hydroxy group, a nitrogen containing 5-6 membered heterocyclic ring which may be unsaturated, partially saturated or wholly saturated and may contain 1 to 3 further heteroatoms selected from N, S and O wherein one of the carbons of the ring may optionally bear a group carboxy, sulfo, carboxy(C₁-C₄)alkyl or sulfo(C₁-C₄)alkyl and the ring nitrogen atom may optionally be substituted by (C₁-C₄)alkyl, or carboxy(C₁-C₄)alkyl, sulfo(C₁-C₄)alkyl, or benzyl;

(C₃-C₆)alkenyl, optionally substituted by carboxy or sulfo;

1-deoxy-1-glucityl;

2-deoxy-2-glucosyl;

a fully saturated 5 to 7 membered nitrogen containing heterocyclic ring wherein the nitrogen atom may be optionally substituted by (C₁-C₄)alkyl or benzyl and one or two carbons of the ring skeleton may bear a substituent selected from (C₁-C₄)alkyl, carboxy and sulfo;

or R₂ and R₃

taken together with the adjacent nitrogen atom represent a fully saturated 5-7 membered heterocyclic ring which may optionally contain a further heteroatom selected from O, S and N, and may optionally bear one or two substituents on the ring carbons selected from (C₁-C₄)alkyl, benzyl, carboxy, sulfo, carboxy(C₁-C₄)alkyl, and sulfo(C₁-C₄)alkyl;

R₄

represents:

hydrogen,

methyl, or

hydroxymethyl;

with the proviso that when R₄ is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and R₁ is methyl;

and the pharmaceutically acceptable addition salts thereof, which comprises reacting an antibiotic GE 2270 compound having formula II:



W represents a carboxy or an activated ester function;

R₁ represents hydrogen or methyl;

R₄ represents hydrogen, methyl or

with the proviso that when R₄ is hydrogen or hydroxymethyl, then simultaneously R is methoxymethyl and R₁ is methyl, with a selected amine of formula HNR₂R₃ wherein R₂ and R₃ have the same meanings as in claim 1, in an inert organic solvent and, when W is carboxy, in the presence of a condensing agent.

12. A process according to claim 11 wherein the condensing agents are selected from (C₁-C₄)alkyl, phenyl or heterocyclic phosphorazidates such as, diphenylphosphorazidate (DPPA), diethylphosphorazidate, di-(4-nitrophenyl)phosphorazidate, dimorpholylphosphorazidate and diphenylphosphorochloridate, or benzotriazol-1-yl-oxy-trispyrrolidinophosphoniumhexafluorophosphate (PyBOP).


13. A process according to claims 11 and 12 wherein the amine reactant HNR_2R_3 is used in a 1 to 2 fold molar excess with respect to the antibiotic starting material and the reaction temperature is comprised between 0 and 20 °C.

- 45 **14.** Use of a compound according to any of claims 1 to 10 for preparing a medicament for use as an antibiotic.

Patentansprüche


Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, MC, NL, SE, PT

1. Amidderivat des Antibiotikums GE 2270 mit folgender Formel I



wobei:

R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe darstellt,
 R₁ ein Wasserstoffatom oder eine Methylgruppe darstellt,
 Y eine Gruppe der Formel




darstellt,

wobei:

R₂ ein Wasserstoffatom, einen (C₁-C₄)-Alkyl-, Amino-(C₂-C₄)-alkyl-, (C₁-C₄)-Alkylamino-(C₁-C₄)-alkyl- oder Di-(C₁-C₄)-alkylamino-(C₁-C₄)-alkylrest darstellt,
 R₃ ein Wasserstoffatom, einen linearen oder verzweigten (C₁-C₁₄)-Alkylrest, der 1 bis 3 Substituenten trägt, ausgewählt aus: einer Carboxy-, Sulfo-, Phosphono-, Aminogruppe, die gegebenenfalls mit einer

Niederalkoxycarbonyl- oder einer Benzyloxycarbonylgruppe geschützt sein kann, einem (C₁-C₄)-Alkylaminorest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einem Di-(C₁-C₄)-alkylamino-, Hydroxyl-, Halogen-, (C₁-C₄)-Alkoxyrest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einem (C₁-C₄)-Alkoxy-carbonyl-, Mercapto-, (C₁-C₄)-Alkylthio-rest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einer Phenylgruppe, die gegebenenfalls mit 1 bis 3 Substituenten substituiert sein kann, ausgewählt aus Carboxy-, Sulfo-, Hydroxyl-, Halogen- und Mercaptogruppen, einem Carbamyl-, (C₁-C₆)-Alkylcarbamylrest, wobei die Alkyleinheit gegebenenfalls mit 1 oder 2 Substituenten, ausgewählt aus Carboxy-, Amino-, (C₁-C₄)-Alkylamino- und Di-(C₁-C₄)-alkylaminoresten substituiert sein kann, einem Di-(C₁-C₄)-alkylcarbamylrest, wobei die Alkyleinheiten zusammen mit dem benachbarten Stickstoffatom auch einen gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen können, der gegebenenfalls mit einer Carboxy- oder Carbamylgruppe an einem der Ringkohlenstoffatome substituiert sein kann und gegebenenfalls eine weitere Heterogruppe, ausgewählt aus O, S und N enthalten kann, einer Benzoylaminogruppe, wobei die Phenylgruppe mit 1 bis 3 Hydroxylgruppen substituiert sein kann, einem stickstoffhaltigen 5-6-gliedrigen heterocyclischen Ring, der ungesättigt, teilweise gesättigt oder vollständig gesättigt sein kann und 1 bis 3 weitere Heteroatome, ausgewählt aus N, S und O enthalten kann, wobei eines der Kohlenstoffatome des Rings gegebenenfalls einen Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- oder Sulfo-(C₁-C₄)-alkylrest tragen kann und das Ringstickstoffatom gegebenenfalls mit einer (C₁-C₄)-Alkyl-, Carboxy-(C₁-C₄)-alkyl-, Sulfo-(C₁-C₄)-alkyl- oder Benzylgruppe substituiert sein kann, einen (C₃-C₆)-Alkenylrest, der gegebenenfalls mit einer Carboxy- oder Sulfogruppe substituiert ist, eine 1-Deoxy-1-glucityl-, eine 2-Deoxy-2-glucosylgruppe, einen vollständig gesättigten 5 bis 7-gliedrigen stickstoffhaltigen heterocyclischen Ring, wobei das Stickstoffatom gegebenenfalls mit einem (C₁-C₄)-Alkylrest oder einer Benzylgruppe substituiert sein kann und ein oder zwei Kohlenstoffatome des Ringskeletts einen Substituenten, ausgewählt aus (C₁-C₄)-Alkylresten, Carboxy- und Sulfogruppen tragen kann, darstellt, oder R₂ und R₃ zusammen mit dem benachbarten Stickstoffatom einen vollständig gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen, der gegebenenfalls ein weiteres Heteroatom, ausgewählt aus O, S und N enthalten kann, und gegebenenfalls ein oder zwei Substituenten, ausgewählt aus (C₁-C₄)-Alkyl-, Benzyl-, Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl und Sulfo-(C₁-C₄)-alkylresten, an den Ringkohlenstoffatomen tragen kann, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, mit der Maßgabe, daß, wenn R₄ ein Wasserstoffatom oder eine Hydroxymethylgruppe darstellt, dann gleichzeitig R eine Methoxymethylgruppe und R₁ eine Methylgruppe ist, und pharmazeutisch verträgliche Säureadditionssalze davon.

2. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe darstellt und die anderen Substituenten die in Anspruch 1 angegebene Bedeutung haben.
3. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe darstellt, R₁ und R₄ eine Methylgruppe darstellen, und Y eine Gruppe der Formel




darstellt, wobei R₂ ein Wasserstoffatom ist und R₃ die in Anspruch 1 angegebene Bedeutung hat.

4. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe ist, R₁ und R₄ eine Methylgruppe darstellen, und Y eine Aminoereinheit ist, die sich von einer natürlichen Aminosäure, wie zum Beispiel Glycin, Ornithin, Serin, Asparaginsäure, Tyrosin, Leucin, Phenylalanin, Methionin, Prolin, Threonin, Lysin oder einem synthetischen Dipeptid, wie Glycyllysin, Serylprolin, Glycylprolinamid, Tyrosylprolinamid, Threonylprolinamid, Leucylprolinamid, ableitet.
5. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe ist, R₁ und R₄ Methylgruppen sind, Y eine Gruppe NR₂R₃ ist, wobei R₂ ein Wasserstoffatom ist und R₃ ein linearer Alkylrest mit

vorzugsweise 3 bis 12 Kohlenstoffatomen, stärker bevorzugt 3 bis 7 Kohlenstoffatomen, ist, der mit einer Gruppe, ausgewählt aus COOH, SO₃H und PO₃H₂ substituiert ist.

6. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe ist, R₁ und R₄ Methylgruppen sind, Y eine Gruppe NR₂R₃ ist, in der R₂ ein Wasserstoffatom ist und R₃ CH₂CH₂CH₂CH₂CH₂-COOH ist.

7. Verbindung nach Anspruch 1, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe darstellt, R₁ ein Wasserstoffatom oder eine Methylgruppe darstellt, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, und Y eine Gruppe der Formel



darstellt, in der R₂ ein Wasserstoffatom ist und R₃ die in Anspruch 1 angegebene Bedeutung hat.

8. Verbindung nach Anspruch 7, wobei Y eine Aminoereinheit ist, die sich von einer natürlichen Aminosäure, wie zum Beispiel Glycin, Ornithin, Serin, Asparaginsäure, Tyrosin, Leucin, Phenylalanin, Methionin, Prolin, Threonin, Lysin, oder einem synthetischen Dipeptid, wie Glycyllysin, Serylprolin, Glycylprolinamid, Tyrosylprolinamid, Threonylprolinamid, Leucylprolinamid, ableitet.

9. Verbindung nach Anspruch 1, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe ist, R₁ ein Wasserstoffatom oder eine Methylgruppe ist, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe ist, und Y eine Gruppe NR₂R₃ ist, in der R₂ ein Wasserstoffatom ist und R₃ ein linearer Alkylrest mit vorzugsweise 3 bis 12 Kohlenstoffatomen, stärker bevorzugt 3 bis 7 Kohlenstoffatomen, ist, der mit einer Gruppe, ausgewählt aus COOH, SO₃H und PO₃H₂, substituiert ist.

10. Verbindung nach Anspruch 1, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe ist, R₁ ein Wasserstoffatom oder eine Methylgruppe ist, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe ist, und Y eine Gruppe NR₂R₃ ist, in der R₂ ein Wasserstoffatom und R₃ CH₂CH₂CH₂CH₂CH₂-COOH ist.

11. Verfahren zur Herstellung einer Verbindung nach Anspruch 1, umfassend die Umsetzung einer Antibiotikum-GE 2270-Verbindung mit Formel II:



15

20

25

30

35

4

4.

5.

- 5




Y eine Gruppe der Formel



50

R₃ ein Wasserstoffatom, einen linearen oder verzweigten (C₁-C₁₄)-Alkylrest, der 1 bis 3 Substituenten trägt, ausgewählt aus: einer Carboxy-, Sulfo-, Phosphono-, Aminogruppe, die gegebenenfalls mit einer Niederalkoxycarbonyl- oder einer Benzyloxycarbonylgruppe geschützt sein kann, einem (C₁-C₄)-Alkylaminorest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einem Di-(C₁-C₄)-alkylamino-, Hydroxyl-, (C₁-C₄)-Alkoxyrest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einem (C₁-C₄)-Alkoxycarbonyl-, Mercapto-, (C₁-C₄)-

Alkylthiorest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einer Phenylgruppe, die gegebenenfalls mit 1 bis 3 Substituenten substituiert sein kann, ausgewählt aus Carboxy-, Sulfo-, Hydroxyl-, Halogen- und Mercaptogruppen, einem Carbamyl-, (C₁-C₆)-Alkylcarbamylrest, wobei die Alkyleinheit gegebenenfalls mit 1 oder 2 Substituenten, ausgewählt aus Carboxy-, Amino-, (C₁-C₄)-Alkylamino- und Di-(C₁-C₄)-alkylaminoesten substituiert sein kann, einem Di-(C₁-C₄)-alkylcarbamylrest, wobei die Alkyleinheiten zusammen mit dem benachbarten Stickstoffatom auch einen gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen können, der gegebenenfalls mit einer Carboxy- oder Carbamylgruppe an einem der Ringkohlenstoffatome substituiert sein kann und gegebenenfalls eine weitere Heterogruppe, ausgewählt aus O, S und N enthalten kann, einer Benzoylamino-
 10 gruppe, wobei die Phenylgruppe mit 1 bis 3 Hydroxylgruppen substituiert sein kann, einem stickstoffhaltigen 5-6-gliedrigen heterocyclischen Ring, der ungesättigt, teilweise gesättigt oder vollständig gesättigt sein kann und 1 bis 3 weitere Heteroatome, ausgewählt aus N, S und O enthalten kann, wobei eines der Kohlenstoffatome des Rings gegebenenfalls einen Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- oder Sulfo-(C₁-C₄)-alkylrest tragen kann und das Ringstickstoffatom gegebenenfalls mit einer (C₁-C₄)-Alkyl-, Carboxy-(C₁-C₄)-alkyl-, Sulfo-(C₁-C₄)-alkyl- oder Benzylgruppe substituiert sein kann, einen (C₃-C₆)-Alkenylrest, der gegebenenfalls mit einer Carboxy- oder Sulfogruppe substituiert ist, eine 1-Deoxy-1-glucityl-, eine 2-Deoxy-2-glucosylgruppe, einen vollständig gesättigten 5 bis 7-gliedrigen stickstoffhaltigen heterocyclischen Ring, wobei das Stickstoffatom gegebenenfalls mit einem (C₁-C₄)-Alkylrest oder einer Benzylgruppe substituiert sein kann und ein oder zwei Kohlenstoffatome des Ringskeletts einen Substituenten, ausgewählt aus (C₁-C₄)-Alkylresten, Carboxy- und Sulfogruppen tragen kann, darstellt,
 20 oder R₂ und R₃ zusammen mit dem benachbarten Stickstoffatom einen vollständig gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen, der gegebenenfalls ein weiteres Heteroatom, ausgewählt aus O, S und N enthalten kann, und gegebenenfalls ein oder zwei Substituenten, ausgewählt aus (C₁-C₄)-Alkyl-, Benzyl-, Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- und Sulfo-(C₁-C₄)-alkylresten, an den Ringkohlenstoffatomen tragen kann,
 25 R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, mit der Maßgabe, daß, wenn R₄ ein Wasserstoffatom oder eine Hydroxymethylgruppe darstellt, dann gleichzeitig R eine Methoxymethylgruppe und R₁ ein Methylgruppe ist, und pharmazeutisch verträglichen Säureadditionssalzen davon, umfassend die Umsetzung einer Antibiotikum-GE 2270-Verbindung mit Formel II:




wobei W eine Carboxygruppe oder eine aktivierte Esterfunktion darstellt,
 R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe darstellt,

R_1 ein Wasserstoffatom oder eine Methylgruppe darstellt,

R_4 ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, mit der Maßgabe, daß, wenn R_4 ein Wasserstoffatom oder eine Hydroxymethylgruppe ist, dann gleichzeitig R eine Methoxymethylgruppe und R_1 eine Methylgruppe ist, mit einem ausgewählten Amin der Formel HNR_2R_3 , wobei R_2 und R_3 die gleichen Bedeutungen wie vorstehend haben, in einem inerten organischen Lösungsmittel, und, wenn W eine Carboxygruppe ist, in Gegenwart eines Kondensationsmittels.

2. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R eine Methoxymethylgruppe darstellt und die anderen Substituenten die in Anspruch 1 angegebene Bedeutung haben.

3. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R eine Methoxymethylgruppe darstellt, R_1 und R_4 eine Methylgruppe darstellen und Y eine Gruppe der Formel




darstellt, wobei R_2 ein Wasserstoffatom ist und R_3 die in Anspruch 1 angegebene Bedeutung hat.

4. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R eine Methoxymethylgruppe ist, R_1 und R_4 eine Methylgruppe darstellen, und Y eine Aminoereinheit ist, die sich von einer natürlichen Aminosäure, wie zum Beispiel Glycin, Ornithin, Serin, Asparaginsäure, Tyrosin, Leucin, Phenylalanin, Methionin, Prolin, Threonin, Lysin oder einem synthetischen Dipeptid, wie Glycyllysin, Serylprolin, Glycylprolinamid, Tyrosylprolinamid, Threonylprolinamid, Leucylprolinamid, ableitet.

5. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R eine Methoxymethylgruppe ist, R_1 und R_4 Methylgruppen sind, Y eine Gruppe NR_2R_3 ist, wobei R_2 ein Wasserstoffatom ist und R_3 ein linearer Alkylrest mit vorzugsweise 3 bis 12 Kohlenstoffatomen, stärker bevorzugt 3 bis 7 Kohlenstoffatomen, ist, der mit einer Gruppe, ausgewählt aus $COOH$, SO_3H und PO_3H_2 , substituiert ist.

6. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R eine Methoxymethylgruppe ist, R_1 und R_4 Methylgruppen sind, Y eine Gruppe NR_2R_3 ist, in der R_2 ein Wasserstoffatom ist und R_3 $CH_2CH_2CH_2CH_2CH_2-COOH$ ist.

7. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe darstellt, R_1 ein Wasserstoffatom oder eine Methylgruppe darstellt, R_4 ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, und Y eine Gruppe der Formel



darstellt, in der R_2 ein Wasserstoffatom ist und R_3 die in Anspruch 1 angegebene Bedeutung hat.

8. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei Y eine Aminoereinheit ist, die sich von einer natürlichen Aminosäure, wie zum Beispiel Glycin, Ornithin, Serin, Asparaginsäure, Tyrosin, Leucin, Phenylalanin, Methionin, Prolin, Threonin, Lysin, oder einem synthetischen Dipeptid, wie Glycyllysin, Serylprolin, Glycylprolinamid, Tyrosylprolinamid, Threonylprolinamid, Leucylprolina-

mid, ableitet.

- 5 9. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe ist, R_1 ein Wasserstoffatom oder eine Methylgruppe ist, R_4 ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe ist, und Y eine Gruppe NR_2R_3 ist, in der R_2 ein Wasserstoffatom ist und R_3 ein linearer Alkylrest mit vorzugsweise 3 bis 12 Kohlenstoffatomen, stärker bevorzugt 3 bis 7 Kohlenstoffatomen, ist, der mit einer Gruppe, ausgewählt aus COOH, SO_3H und PO_3H_2 , substituiert ist.
- 10 10. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung der Formel I, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe ist, R_1 ein Wasserstoffatom oder eine Methylgruppe ist, R_4 ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe ist, und Y eine Gruppe NR_2R_3 ist, in der R_2 ein Wasserstoffatom und R_3 $CH_2CH_2CH_2CH_2CH_2-COOH$ ist.
- 15 11. Verfahren nach einem der Ansprüche 1 bis 10, wobei die Kondensationsmittel ausgewählt sind aus (C_1-C_4)-Alkyl-, Phenyl- oder heterocyclischen Phosphorazidaten, wie Diphenylphosphorazidat (DPPA), Diethylphosphorazidat, Di-(4-nitrophenyl)phosphorazidat, Dimorpholyphosphorazidat und Diphenylphosphorchloridat, oder Benzotriazol-1-yl-oxy-trispyrrolidinophosphoniumhexafluorophosphat (PyBOP).
- 20 12. Verfahren nach einem der Ansprüche 1 bis 11, wobei der Aminumsetzungsteilnehmer HNR_2R_3 in einem 1- bis 2-fachen molaren Überschuß in bezug auf die antibiotische Ausgangssubstanz verwendet wird, und die Umsetzungstemperatur zwischen 0 und 20 °C beträgt.
- 25 13. Verwendung einer nach einem der Ansprüche 1 bis 12 hergestellten Verbindung zur Herstellung eines Medikaments zur Anwendung als Antibiotikum.

30

35


40

45


50

55

1. Amidderivat des Antibiotikums GE 2270 mit folgender Formel I




Y eine Gruppe der Formel



R₃ ein Wasserstoffatom, einen linearen oder verzweigten (C₁-C₄)-Alkylrest, der 1 bis 3 Substituenten trägt, ausgewählt aus: einer Carboxy-, Sulfo-, Phosphono-, Aminogruppe, die gegebenenfalls mit einer Niederalkoxycarbonyl- oder einer Benzoyloxycarbonylgruppe geschützt sein kann, einem (C₁-C₄)-Alkylaminorest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einem Di-(C₁-C₄)-alkylamino-, Hydroxyl-, Halogen-, (C₁-C₄)-Alkoxyrest, wobei die Alkyleinheit gegebenenfalls

mit einer Carboxygruppe substituiert sein kann, einem (C₁-C₄)-Alkoxy-carbonyl-, Mercapto-, (C₁-C₄)-Alkylthio-, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einer Phenylgruppe, die gegebenenfalls mit 1 bis 3 Substituenten substituiert sein kann, ausgewählt aus Carboxy-, Sulfo-, Hydroxyl-, Halogen- und Mercapto-gruppen, einem Carbamyl-, (C₁-C₆)-Alkylcarbamylrest, wobei die Alkyleinheit gegebenenfalls mit 1 oder 2 Substituenten, ausgewählt aus Carboxy-, Amino-, (C₁-C₄)-Alkylamino- und Di-(C₁-C₄)-alkylamino-resten substituiert sein kann, einem Di-(C₁-C₄)-alkylcarbamylrest, wobei die Alkyleinheiten zusammen mit dem benachbarten Stickstoffatom auch einen gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen können, der gegebenenfalls mit einer Carboxy- oder Carbamylgruppe an einem der Ringkohlenstoffatome substituiert sein kann und gegebenenfalls eine weitere Heterogruppe, ausgewählt aus O, S und N enthalten kann, einer Benzoylamino-gruppe, wobei die Phenylgruppe mit 1 bis 3 Hydroxylgruppen substituiert sein kann, einem stickstoffhaltigen 5-6-gliedrigen heterocyclischen Ring, der ungesättigt, teilweise gesättigt oder vollständig gesättigt sein kann und 1 bis 3 weitere Heteroatome, ausgewählt aus N, S und O enthalten kann, wobei eines der Kohlenstoffatome des Rings gegebenenfalls einen Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- oder Sulfo-(C₁-C₄)-alkylrest tragen kann, und das Ringstickstoffatom gegebenenfalls mit einer (C₁-C₄)-Alkyl-, Carboxy-(C₁-C₄)-alkyl-, Sulfo-(C₁-C₄)-alkyl- oder Benzylgruppe substituiert sein kann, einen (C₃-C₆)-Alkenylrest, der gegebenenfalls mit einer Carboxy- oder Sulfogruppe substituiert ist, eine 1-Deoxy-1-glucityl-, eine 2-Deoxy-2-glucosylgruppe, einen vollständig gesättigten 5 bis 7-gliedrigen stickstoffhaltigen heterocyclischen Ring, wobei das Stickstoffatom gegebenenfalls mit einem (C₁-C₄)-Alkylrest oder einer Benzylgruppe substituiert sein kann und ein oder zwei Kohlenstoffatome des Ringskeletts einen Substituenten, ausgewählt aus (C₁-C₄)-Alkylresten, Carboxy- und Sulfogruppen, tragen kann, darstellt, oder R₂ und R₃ zusammen mit dem benachbarten Stickstoffatom einen vollständig gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen, der gegebenenfalls ein weiteres Heteroatom, ausgewählt aus O, S und N, enthalten kann, und gegebenenfalls ein oder zwei Substituenten, ausgewählt aus (C₁-C₄)-Alkyl-, Benzyl-, Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- und Sulfo-(C₁-C₄)-alkylresten, an den Ringkohlenstoffatomen tragen kann, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, mit der Maßgabe, daß, wenn R₄ ein Wasserstoffatom oder eine Hydroxymethylgruppe darstellt, dann gleichzeitig R eine Methoxymethylgruppe und R₁ ein Methylgruppe ist, und pharmazeutisch verträgliche Säureadditionssalze davon.


2. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe darstellt und die anderen Substituenten die in Anspruch 1 angegebene Bedeutung haben.
3. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe darstellt, R₁ und R₄ eine Methylgruppe darstellen, und Y eine Gruppe der Formel





darstellt, wobei R₂ ein Wasserstoffatom ist und R₃ die in Anspruch 1 angegebene Bedeutung hat.

4. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe ist, R₁ und R₄ eine Methylgruppe darstellen, und Y eine Aminosäureeinheit ist, die sich von einer natürlichen Aminosäure, wie zum Beispiel Glycin, Ornithin, Serin, Asparaginsäure, Tyrosin, Leucin, Phenylalanin, Methionin, Prolin, Threonin, Lysin oder einem synthetischen Dipeptid, wie Glycyl-Lysin, Serylprolin, Glycylprolinamid, Tyrosylprolinamid, Threonylprolinamid, Leucylprolinamid, ableitet.
5. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe ist, R₁ und R₄ Methylgruppen sind, Y eine Gruppe NR₂R₃ ist, wobei R₂ ein Wasserstoffatom ist und R₃ ein linearer Alkylrest mit vorzugsweise 3 bis 12 Kohlenstoffatomen, stärker bevorzugt 3 bis 7 Kohlenstoffatomen, darstellt, der mit einer Gruppe, ausgewählt aus COOH, SO₃H und PO₃H₂, substituiert ist.

6. Verbindung nach Anspruch 1, wobei R eine Methoxymethylgruppe ist, R₁ und R₄ Methylgruppen sind, Y eine Gruppe NR₂R₃ ist, in der R₂ ein Wasserstoffatom ist und R₃ CH₂CH₂CH₂CH₂CH₂-COOH ist.
7. Verbindung nach Anspruch 1, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe darstellt, R₁ ein Wasserstoffatom oder eine Methylgruppe darstellt, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, und Y eine Gruppe der Formel



- darstellt, in der R₂ ein Wasserstoffatom ist und R₃ die in Anspruch 1 angegebene Bedeutung hat.
8. Verbindung nach Anspruch 7, wobei Y eine Aminoereinheit ist, die sich von einer natürlichen Aminosäure, wie zum Beispiel Glycin, Ornithin, Serin, Asparaginsäure, Tyrosin, Leucin, Phenylalanin, Methionin, Prolin, Threonin, Lysin, oder einem synthetischen Dipeptid, wie Glycyllysin, Serylprolin, Glycylprolinamid, Tyrosylprolinamid, Threonylprolinamid, Leucylprolinamid, ableitet.
9. Verbindung nach Anspruch 1, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe ist, R₁ ein Wasserstoffatom oder eine Methylgruppe ist, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe ist, und Y eine Gruppe NR₂R₃ ist, in der R₂ ein Wasserstoffatom ist, und R₃ ein linearer Alkylrest mit vorzugsweise 3 bis 12 Kohlenstoffatomen, stärker bevorzugt 3 bis 7 Kohlenstoffatomen, ist, der mit einer Gruppe, ausgewählt aus COOH, SO₃H und PO₃H₂, substituiert ist.
10. Verbindung nach Anspruch 1, wobei R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe ist, R₁ ein Wasserstoffatom oder eine Methylgruppe ist, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe ist, und Y eine Gruppe NR₂R₃ ist, in der R₂ ein Wasserstoffatom und R₃ CH₂CH₂CH₂CH₂CH₂-COOH ist.




50 R₂ ein Wasserstoffatom, einen (C₁-C₄)-Alkyl-, Amino-(C₂-C₄)-alkyl-, (C₁-C₄)-Alkylamino-(C₁-C₄)-alkyl- oder Di-(C₁-C₄)-alkylamino-(C₁-C₄)-alkylrest darstellt,
R₃ ein Wasserstoffatom, einen linearen oder verzweigten (C₁-C₁₄)-Alkylrest, der 1 bis 3 Substituenten trägt, ausgewählt aus: einer Carboxy-, Sulfo-, Phosphono-, Aminogruppe, die gegebenenfalls mit einer Niederalkoxycarbonyl- oder einer Benzoyloxycarbonylgruppe geschützt sein kann, einem (C₁-C₄)-Alkylamino-
55 rest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxylgruppe substituiert sein kann, einem Di-(C₁-C₄)-alkylamino-, Hydroxyl-, Halogen-, (C₁-C₄)-Alkoxyrest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, einem (C₁-C₄)-Alkoxy-carbonyl-, Mercapto-, (C₁-C₄)-Alkylthio-
rest, wobei die Alkyleinheit gegebenenfalls mit einer Carboxygruppe substituiert sein kann, eine

Phenylgruppe, die gegebenenfalls mit 1 bis 3 Substituenten substituiert sein kann, ausgewählt aus Carboxy-, Sulfo-, Hydroxyl-, Halogen- und Mercaptogruppen, einem Carbamyl-, (C₁-C₆)-Alkylcarbamylrest, wobei die Alkyleinheit gegebenenfalls mit 1 oder 2 Substituenten, ausgewählt aus Carboxy-, Amino-, (C₁-C₄)-Alkylamino- und Di-(C₁-C₄)-alkylaminoresten substituiert sein kann, einem Di-(C₁-C₄)-alkylcarbamylrest, wobei die Alkyleinheiten zusammen mit dem benachbarten Stickstoffatom auch einen gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen können, der gegebenenfalls mit einer Carboxy- oder Carbamylgruppe an einem der Ringkohlenstoffatome substituiert sein kann und gegebenenfalls eine weitere Heterogruppe, ausgewählt aus O, S und N enthalten kann, einer Benzoylamino-
 10 gruppe, wobei die Phenylgruppe mit 1 bis 3 Hydroxylgruppen substituiert sein kann, einem stickstoffhaltigen 5-6-gliedrigen heterocyclischen Ring, der ungesättigt, teilweise gesättigt oder vollständig gesättigt sein kann und 1 bis 3 weitere Heteroatome, ausgewählt aus N, S und O, enthalten kann, wobei eines der Kohlenstoffatome des Rings gegebenenfalls einen Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- oder Sulfo-(C₁-C₄)-alkylrest tragen kann und das Ringstickstoffatom gegebenenfalls mit einer (C₁-C₄)-Alkyl-, Carboxy-(C₁-C₄)-alkyl-, Sulfo-(C₁-C₄)-alkyl- oder Benzylgruppe substituiert sein kann, einen (C₃-C₆)-Alkenylrest, der gegebenenfalls mit einer Carboxy- oder Sulfogruppe substituiert ist, eine 1-Deoxy-1-glucityl-, eine 2-Deoxy-2-glucosylgruppe, einen vollständig gesättigten 5 bis 7-gliedrigen stickstoffhaltigen heterocyclischen Ring, wobei das Stickstoffatom gegebenenfalls mit einem (C₁-C₄)-Alkylrest oder einer Benzylgruppe substituiert sein kann und ein oder zwei Kohlenstoffatome des Ringskeletts einen Substituenten, ausgewählt aus (C₁-C₄)-Alkylresten, Carboxy- und Sulfogruppen, tragen kann, darstellt,

oder R₂ und R₃ zusammen mit dem benachbarten Stickstoffatom einen vollständig gesättigten 5-7-gliedrigen heterocyclischen Ring darstellen, der gegebenenfalls ein weiteres Heteroatom, ausgewählt aus O, S und N, enthalten kann, und gegebenenfalls ein oder zwei Substituenten, ausgewählt aus (C₁-C₄)-Alkyl-, Benzyl-, Carboxy-, Sulfo-, Carboxy-(C₁-C₄)-alkyl- und Sulfo-(C₁-C₄)-alkylresten, tragen kann, R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt,


mit der Maßgabe, daß, wenn R₄ ein Wasserstoffatom oder eine Hydroxymethylgruppe darstellt, dann gleichzeitig R eine Methoxymethylgruppe und R₁ ein Methylgruppe ist, und pharmazeutisch verträglichen Säureadditionssalzen davon, umfassend die Umsetzung einer Antibiotikum-GE 2270-Verbindung der Formel II:




wobei W eine Carboxygruppe oder eine aktivierte Esterfunktion darstellt,
 R ein Wasserstoffatom, eine Hydroxymethyl- oder Methoxymethylgruppe darstellt,

R₄ ein Wasserstoffatom, eine Methyl- oder Hydroxymethylgruppe darstellt, mit der Maßgabe, daß, wenn R₄ ein Wasserstoffatom oder eine Hydroxymethylgruppe ist, dann gleichzeitig R eine Methoxymethylgruppe und R₁ eine Methylgruppe ist, mit einem ausgewählten Amin der Formel HNR₂R₃, wobei R₂

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, MC, NL, SE, PT



le groupe hydroxyméthyle ou
 le groupe méthoxyméthyle;
 R₁ représente un atome d'hydrogène ou le groupe méthyle;
 Y représente un groupe de formule



dans laquelle

R₂ représente un atome d'hydrogène,
 un groupe alkyle en C₁-C₄, un groupe aminoalkyle(C₂-C₄),
 un groupe alkyl(C₁-C₄)amino-alkyle(C₁-C₄) ou
 un groupe dialkyl(C₁-C₄)amino-alkyle(C₁-C₄);

R₃ représente un atome d'hydrogène,
 un groupe alkyle en C₁-C₁₄ linéaire ou ramifié portant 1 à 3 substituants choisis parmi
 des groupes carboxy, sulfo, phosphono, amino qui peut être éventuellement protégé
 par un groupe(alcoxy inférieur)carbonyle ou benzyloxycarbonyl, alkyl(C₁-C₄)amino
 dans lequel le fragment alkyle peut être éventuellement substitué par un groupe
 carboxy, dialkyl(C₁-C₄)amino, hydroxy, halogéno, alcoxy en C₁-C₄ dans lequel le
 fragment alkyle peut être éventuellement substitué par un groupe carboxy, alcoxy(C₁-
 C₄)carbonyle, mercapto, alkyl(C₁-C₄)thio dans lequel le fragment alkyle peut être
 éventuellement substitué par un groupe carboxy, phényle qui peut être éventuellement
 substitué par 1 à 3 substituants choisis parmi des groupes carboxy, sulfo, hydroxy
 halogéno

et mercapto, carbamoyl, alkyl(C₁-C₆)carbamoyl dans lequel le fragment alkyle peut
 être éventuellement substitué par 1 ou 2 substituants choisis parmi des groupes
 carboxy, amino, alkyl(C₁-C₄)amino et dialkyl(C₁-C₄)amino, dialkyl(C₁-C₄)carbamoyl
 dans lequel les fragments alkyle conjointement avec l'atome d'azote adjacent peuvent
 également représenter un cycle hétérocyclique saturé à 5-7 chaînons qui peut éven-
 tuellement être substitué par un groupe carboxy ou carbamoyl sur l'un des atomes de
 carbone formant le cycle, et qui peut éventuellement contenir un autre hétéroatome
 choisi parmi O, S et N, benzoylamino dans lequel le fragment phényle peut être
 substitué par 1 à 3 groupes hydroxy, un cycle hétérocyclique azoté contenant 5-6
 chaînons, qui peut être insaturé, partiellement saturé ou totalement saturé et peut
 contenir 1 à 3 autres hétéroatomes choisis parmi N, S et O, l'un des atomes de
 carbone du cycle pouvant éventuellement porter un groupe carboxy, sulfo, carboxy-
 alkyle(C₁-C₄) ou sulfo-alkyle(C₁-C₄) et l'atome d'azote du cycle pouvant éventuellement
 être substitué par un groupe alkyle en C₁-C₄, carboxy-alkyle(C₁-C₄), sulfo-alkyle(C₁-C₄)
 ou benzyle;

un groupe alcényle en C₃-C₆, éventuellement substitué par le groupe carboxy ou sulfo;
 le groupe 1-désoxy-1-glucityl;

le groupe 2-désoxy-2-glucosyl;

un cycle hétérocyclique azoté à 5-7 chaînons, totalement saturé, dans lequel l'atome
 d'azote peut être éventuellement substitué par un groupe alkyle en C₁-C₄ ou benzyle,
 et 1 ou 2 atomes de carbone du squelette cyclique peuvent porter un substituant choisi
 parmi des groupes alkyle en C₁-C₄, carboxy et sulfo;


ou

R₂ et R₃ forment ensemble, avec l'atome d'azote adjacent, un cycle hétérocyclique à 5-7
 chaînons totalement saturé, qui peut éventuellement contenir un autre hétéroatome
 choisi parmi O, S et N, et peut éventuellement porter 1 ou 2 substituants sur les
 atomes de carbone formant le cycle, choisis parmi des groupes alkyle en C₁-C₄,
 benzyle, carboxy, sulfo, carboxy-alkyle(C₁-C₄) et sulfo-alkyle(C₁-C₄);

R₄ représente un atome d'hydrogène,
 le groupe méthyle ou


le groupe hydroxyméthyle,
 sous réserve que lorsque R_4 est un atome d'hydrogène ou le groupe hydroxyméthyle, alors R est
 simultanément le groupe méthoxyméthyle et R_1 est le groupe méthyle;
 et sels d'addition pharmaceutiquement acceptables de celui-ci.

- 5
2. Composé selon la revendication 1, dans lequel R représente le groupe méthoxyméthyle et les autres substituants sont tels que définis dans la revendication 1.
3. Composé selon la revendication 1, dans lequel R représente le groupe méthoxyméthyle, R_1 et R_4
 10 représentent chacun le groupe méthyle et Y représente un groupe de formule



dans laquelle R_2 est un atome d'hydrogène et R_3 est tel que défini dans la revendication 1.

- 20 4. Composé selon la revendication 1, dans lequel R est le groupe méthoxyméthyle, R_1 et R_4 sont chacun le groupe méthyle et Y est un fragment aminé qui dérive d'un aminoacide naturel tel que la glycine, l'ornithine, la sérine, l'acide aspartique, la tyrosine, la leucine, la phénylalanine, la méthionine, la proline, la thréonine, la lysine ou d'un dipeptide synthétique tel que la glycyllysine, la sérylproline, le glycyprolinamide, le tyrosylprolinamide, le thréonylprolinamide, le leucylprolinamide.
- 25 5. Composé selon la revendication 1, dans lequel R est le groupe méthoxyméthyle, R_1 et R_4 sont chacun le groupe méthyle, Y est un groupe NR_2R_3 dans lequel R_2 est un atome d'hydrogène et R_3 est une chaîne alkyle linéaire ayant de préférence de 3 à 12 atomes de carbone, encore mieux de 3 à 7 atomes de carbone, substituée par un groupe choisi parmi $COOH$, SO_3H et PO_3H_2 .
- 30 6. Composé selon la revendication 1, dans lequel R est le groupe méthoxyméthyle, R_1 et R_4 sont chacun le groupe méthyle, Y est un groupe NR_2R_3 dans lequel R_2 est un atome d'hydrogène, et R_3 est $CH_2CH_2CH_2CH_2CH_2-COOH$.
- 35 7. Composé selon la revendication 1, dans lequel R représente un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R_1 représente un atome d'hydrogène ou le groupe méthyle, R_4 représente un atome d'hydrogène, le groupe méthyle ou hydroxyméthyle, et Y représente un groupe de formule




dans laquelle R_2 est un atome d'hydrogène et R_3 est tel que défini dans la revendication 1.

- 50 8. Composé selon la revendication 7, dans lequel Y est un fragment aminé qui dérive d'un aminoacide naturel tel que la glycine, l'ornithine, la sérine, l'acide aspartique, la tyrosine, la leucine, la phénylalanine, la méthionine, la proline, la thréonine, la lysine, ou d'un dipeptide synthétique tel que la glycyllysine, la sérylproline, le glycyprolinamide, le tyrosylprolinamide, le thréonylprolinamide, le leucylprolinamide.
- 55 9. Composé selon la revendication 1, dans lequel R est un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R_1 est un atome d'hydrogène ou le groupe méthyle, R_4 est un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle, et Y est un groupe NR_2R_3 dans lequel R_2 est un atome d'hydrogène et R_3 est une chaîne alkyle linéaire ayant de préférence 3 à 12 atomes de

carbone, encore mieux de 3 à 7 atomes de carbone, substituée par un groupe choisi parmi COOH, SO₃H et PO₃H₂.

10. Composé selon la revendication 1, dans lequel R est un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ est un atome d'hydrogène ou le groupe méthyle, R₄ est un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle, et Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est CH₂CH₂CH₂CH₂CH₂-COOH.
11. Procédé pour la préparation d'un composé selon la revendication 1, qui comprend la mise en réaction d'un composé antibiotique GE 2270 de formule II:



dans laquelle

- W représente le groupe carboxy ou une fonction ester activée;
 R représente un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle;
 R₁ représente un atome d'hydrogène ou le groupe méthyle;
 R₄ représente un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle;
 sous réserve que lorsque R₄ est un atome d'hydrogène ou le groupe hydroxyméthyle, R est alors simultanément le groupe méthoxyméthyle et R₁ est le groupe méthyle,
 avec une amine choisie de formule HNR₂R₃ dans laquelle R₂ et R₃ ont les mêmes significations que dans la revendication 1, dans un solvant organique inerte, et, lorsque W est le groupe carboxy, en présence d'un agent de condensation.

12. Procédé selon la revendication 11, dans lequel les agents de condensation sont choisis parmi des alkyl-(C₁-C₄)-, phénylphosphorazidates ou phosphorazidates hétérocycliques tels que le diphénylphosphorazidate (DPPA), le diéthylphosphorazidate, le di-(4-nitrophényl)phosphorazidate, le dimorpholylphosphorazidate et le diphénylphosphorochloridate ou le benzotriazol-1-yl-oxyltrispyrrolidinophosphonium hexafluorophosphate (PyBOP).

13. Procédé selon les revendications 11 et 12, dans lequel le partenaire réactionnel aminé HNR₂R₃ est utilisé en un excès molaire de 1 à 2 fois par rapport au produit de départ antibiotique, et la température de la réaction est comprise entre 0 et 20 °C.

14. Composé selon l'une quelconque des revendications 1 à 10, pour utilisation en tant que médicament.

15. Composition pharmaceutique contenant un composé selon l'une quelconque des revendications 1 à 10, en tant que composant actif en mélange avec un véhicule pharmaceutiquement acceptable.

16. Utilisation d'un composé selon l'une quelconque des revendications 1 à 10, pour la fabrication d'un médicament pour utilisation en tant qu'antibiotique.

Revendications pour l'Etat contractant suivant : ES

1. Procédé pour la préparation d'un dérivé de type amide de l'antibiotique GE 2270 de formule I suivante

10

15

20

25

30

35

40


dans laquelle

R représente un atome d'hydrogène,
le groupe hydroxyméthyle ou
le groupe méthoxyméthyle;

R₁ représente un atome d'hydrogène ou le groupe méthyle;

Y représente un groupe de formule

50



55

R₂ dans laquelle
représente un atome d'hydrogène, un groupe alkyle en C₁-C₄,
un groupe aminoalkyle(C₂-C₄),
un groupe alkyl(C₁-C₄)amino-alkyle(C₁-C₄) ou

R₃un groupe dialkyl(C₁-C₄)amino-alkyle(C₁-C₄);

représente un atome d'hydrogène,

un groupe alkyle en C₁-C₄ linéaire ou ramifié portant 1 à 3 substituants choisis parmi des groupes carboxy, sulfo, phosphono, amino qui peut être éventuellement protégé par un groupe (alcoxy inférieur)carbonyle ou benzyloxycarbonyle, alkyl(C₁-C₄)amino dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, dialkyl(C₁-C₄)amino, hydroxy, halogéno, alcoxy en C₁-C₄ dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, alcoxy(C₁-C₄)carbonyle, mercapto, alkyl(C₁-C₄)thio dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, phényle qui peut être éventuellement substitué par 1 à 3 substituants choisis parmi des groupes carboxy, sulfo, hydroxy, halogéno

et mercapto, carbamoyle, alkyl(C₁-C₆)carbamoyle dans lequel le fragment alkyle peut être éventuellement substitué par 1 ou 2 substituants choisis parmi des groupes carboxy, amino, alkyl(C₁-C₄)amino et dialkyl(C₁-C₄)amino, dialkyl(C₁-C₄)carbamoyle dans lequel les fragments alkyle conjointement avec l'atome d'azote adjacent peuvent également représenter un cycle hétérocyclique saturé à 5-7 chaînons qui peut éventuellement être substitué par un groupe carboxy ou carbamoyle sur l'un des atomes de carbone formant le cycle, et qui peut éventuellement contenir un autre hétéroatome choisi parmi O, S et N, benzoylamino dans lequel le fragment phényle peut être substitué par 1 à 3 groupes hydroxy, un cycle hétérocyclique azoté contenant 5-6 chaînons, qui peut être insaturé, partiellement saturé ou totalement saturé et peut contenir 1 à 3 autres hétéroatomes choisis parmi N, S et O, l'un des atomes de carbone du cycle pouvant éventuellement porter un groupe carboxy, sulfo, carboxy-alkyle(C₁-C₄) ou sulfo-alkyle(C₁-C₄) et l'atome d'azote du cycle pouvant éventuellement être substitué par un groupe alkyle en C₁-C₄, carboxy-alkyle(C₁-C₄), sulfo-alkyle(C₁-C₄) ou benzyle;

un groupe alcényle en C₃-C₆, éventuellement substitué par le groupe carboxy ou sulfo; le groupe 1-désoxy-1-glucityle;

le groupe 2-désoxy-2-glucosyle;

un cycle hétérocyclique azoté à 5-7 chaînons, totalement saturé, dans lequel l'atome d'azote peut être éventuellement substitué par un groupe alkyle en C₁-C₄ ou benzyle, et 1 ou 2 atomes de carbone du squelette cyclique peuvent porter un substituant choisi parmi des groupes alkyle en C₁-C₄, carboxy et sulfo;

ou

R₂ et R₃

forment ensemble, avec l'atome d'azote adjacent, un cycle hétérocyclique à 5-7 chaînons totalement saturé, qui peut éventuellement contenir un autre hétéroatome choisi parmi O, S et N, et peut éventuellement porter 1 ou 2 substituants sur les atomes de carbone formant le cycle, choisis parmi des groupes alkyle en C₁-C₄, benzyle, carboxy, sulfo, carboxy-alkyle(C₁-C₄) et sulfo-alkyle(C₁-C₄);

R₄


représente un atome d'hydrogène,

le groupe méthyle ou

le groupe hydroxyméthyle,

sous réserve que lorsque R₄ est un atome d'hydrogène ou le groupe hydroxyméthyle, R est alors simultanément le groupe méthoxyméthyle et R₁ est le groupe méthyle;

et de ses sels d'addition pharmaceutiquement acceptables, qui comprend la mise en réaction d'un composé antibiotique GE 2270 de formule II:



dans laquelle

W représente le groupe carboxy ou une fonction ester activée;

R représente un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle;

R₁ représente un atome d'hydrogène ou le groupe méthyle;


R₄ représente un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle;

sous réserve que lorsque R₄ est un atome d'hydrogène ou le groupe hydroxyméthyle, R est alors simultanément le groupe méthoxyméthyle et R₁ est le groupe méthyle,

avec une amine choisie de formule HNR₂R₃ dans laquelle R₂ et R₃ ont les mêmes significations que ci-dessus, dans un solvant organique inerte, et, lorsque W est le groupe carboxy, en présence d'un agent de condensation.

2. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R représente le groupe méthoxyméthyle et les autres substituants sont tels que définis dans la revendication 1.

3. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R représente le groupe méthoxyméthyle, R₁ et R₄ représentent chacun le groupe méthyle et Y représente un groupe de formule




dans laquelle R₂ est un atome d'hydrogène et R₃ est tel que défini dans la revendication 1.

4. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R est le groupe méthoxyméthyle, R₁ et R₄ sont chacun le groupe méthyle et Y est un fragment aminé qui dérive d'un aminoacide naturel tel que la glycine, l'ornithine, la sérine, l'acide aspartique, la tyrosine, la leucine, la phénylalanine, la méthionine, la proline, la thréonine, la lysine, ou d'un dipeptide synthétique tel que la glycyllysine, la sérylproline, le glycyprolinamide, le tyrosylprolinamide, le thréonylprolinamide, le leucylprolinamide.

5. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R est le groupe méthoxyméthyle, R₁ et R₄ sont chacun le groupe méthyle, Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est une chaîne alkyle linéaire ayant de préférence de 3 à 12 atomes de carbone, encore mieux de 3 à 7 atomes de carbone, substituée par un groupe choisi parmi COOH, SO₃H et PO₃H₂.

6. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R est le groupe méthoxyméthyle, R₁ et R₄ sont chacun le groupe méthyle, Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène, et R₃ est CH₂CH₂CH₂CH₂CH₂-COOH.

7. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R représente un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ représente un atome d'hydrogène ou le groupe méthyle, R₄ représente un atome d'hydrogène, le groupe méthyle ou hydroxyméthyle, et Y représente un groupe de formule



dans laquelle R₂ est un atome d'hydrogène et R₃ est tel que défini dans la revendication 1.

8. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle Y est un fragment aminé qui dérive d'un aminoacide naturel tel que la glycine, l'ornithine, la sérine, l'acide aspartique, la tyrosine, la leucine, la phénylalanine, la méthionine, la proline, la thréonine, la lysine, ou d'un dipeptide synthétique tel que la glycyllysine, la sérylproline, le glycyprolinamide, le tyrosylprolinamide, le thréonylprolinamide, le leucylprolinamide.

9. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R est un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ est un atome d'hydrogène ou le groupe méthyle, R₄ est un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle, et Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est une chaîne alkyle linéaire ayant de préférence 3 à 12 atomes de carbone, encore mieux de 3 à 7 atomes de carbone, substituée par un groupe choisi parmi COOH, SO₃H et PO₃H₂.

10. Procédé selon la revendication 1, pour la préparation d'un composé de formule I dans laquelle R est un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ est un atome d'hydrogène ou le groupe méthyle, R₄ est un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle, et Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est CH₂CH₂CH₂CH₂CH₂-COOH.

11. Procédé selon l'une quelconque des revendications 1 à 10, dans lequel les agents de condensation sont choisis parmi des alkyl(C₁-C₄)-, phénylphosphorazidates ou phosphorazidates hétérocycliques tels que le diphenylphosphorazidate (DPPA), le diéthylphosphorazidate, le di-(4-nitrophényl)-phosphorazidate, le dimorpholyphosphorazidate et le diphenylphosphorochloridate, ou le benzotriazol-1-yl-oxytris-pyrrolidinophosphonium hexafluorophosphate (Py BOP).

12. Procédé selon l'une quelconque des revendications 1 à 11, dans lequel le partenaire réactionnel aminé HNR₂R₃ est utilisé en un excès molaire de 1 à 2 fois par rapport au produit de départ antibiotique, et la température de la réaction est comprise entre 0 et 20 °C.

13. Utilisation d'un composé préparé selon l'une quelconque des revendications 1 à 12, pour la fabrication d'un médicament pour utilisation en tant qu'antibiotique.

Revendications pour l'Etat contractant suivant : GR

1. Dérivé de type amide de l'antibiotique GE 2270 de formule I suivante

5

10

15

20

25

30

35

dans laquelle

R

représente un atome d'hydrogène,
le groupe hydroxyméthyle ou
le groupe méthoxyméthyle;

40


R₁

représente un atome d'hydrogène ou le groupe méthyle;

Y

représente un groupe de formule .

45



50

R₂dans laquelle
représente un atome d'hydrogène,
un groupe alkyle en C₁-C₄,
un groupe aminoalkyle(C₂-C₄),
un groupe alkyl(C₁-C₄)amino-alkyle(C₁-C₄) ou
un groupe dialkyl(C₁-C₄)amino-alkyle(C₁-C₄);

55

R₃représente un atome d'hydrogène,
un groupe alkyle en C₁-C₁₄ linéaire ou ramifié portant 1 à 3 substituants choisis parmi
des groupes carboxy, sulfo, phosphono, amino qui peut être éventuellement protégé

par un groupe(alcoxy inférieur)carbonyle ou benzyloxycarbonyle, alkyl(C₁-C₄)amino dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, dialkyl(C₁-C₄)amino, hydroxy, halogéno, alcoxy en C₁-C₄ dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, alcoxy(C₁-C₄)carbonyle, mercapto, alkyl(C₁-C₄)thio dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, phényle qui peut être éventuellement substitué par 1 à 3 substituants choisis parmi des groupes carboxy, sulfo, hydroxy, halogéno

et mercapto, carbamoyle, alkyl(C₁-C₆)carbamoyle dans lequel le fragment alkyle peut être éventuellement substitué par 1 ou 2 substituants choisis parmi des groupes carboxy, amino, alkyl(C₁-C₄)amino et dialkyl(C₁-C₄)amino, dialkyl(C₁-C₄)carbamoyle dans lequel les fragments alkyle conjointement avec l'atome d'azote adjacent peuvent également représenter un cycle hétérocyclique saturé à 5-7 chaînons qui peut éventuellement être substitué par un groupe carboxy ou carbamoyle sur l'un des atomes de carbone formant le cycle, et qui peut éventuellement contenir un autre hétéroatome choisi parmi O, S et N, benzoylamino dans lequel le fragment phényle peut être substitué par 1 à 3 groupes hydroxy, un cycle hétérocyclique azoté contenant 5-6 chaînons, qui peut être insaturé, partiellement saturé ou totalement saturé et peut contenir 1 à 3 autres hétéroatomes choisis parmi N, S et O, l'un des atomes de carbone du cycle pouvant éventuellement porter un groupe carboxy, sulfo, carboxy-alkyle(C₁-C₄) ou sulfo-alkyle(C₁-C₄) et l'atome d'azote du cycle pouvant éventuellement être substitué par un groupe alkyle en C₁-C₄, carboxy-alkyle(C₁-C₄), sulfo-alkyle(C₁-C₄) ou benzyle;

un groupe alcényle en C₃-C₆, éventuellement substitué par le groupe carboxy ou sulfo;

le groupe 1-désoxy-1-glucityle;

le groupe 2-désoxy-2-glucosyle;

un cycle hétérocyclique azoté à 5-7 chaînons, totalement saturé, dans lequel l'atome d'azote peut être éventuellement substitué par un groupe alkyle en C₁-C₄ ou benzyle, et 1 ou 2 atomes de carbone du squelette cyclique peuvent porter un substituant choisi parmi des groupes alkyle en C₁-C₄, carboxy et sulfo;

ou

R₂ et R₃ forment ensemble, avec l'atome d'azote adjacent, un cycle hétérocyclique à 5-7 chaînons totalement saturé, qui peut éventuellement contenir un autre hétéroatome choisi parmi O, S et N, et peut éventuellement porter 1 ou 2 substituants sur les atomes de carbone formant le cycle, choisis parmi des groupes alkyle en C₁-C₄, benzyle, carboxy, sulfo, carboxy-alkyle(C₁-C₄) et sulfo-alkyle(C₁-C₄);

R₄ représente un atome d'hydrogène,


le groupe méthyle ou

le groupe hydroxyméthyle,

sous réserve que lorsque R₄ est un atome d'hydrogène ou le groupe hydroxyméthyle, R est alors simultanément le groupe méthoxyméthyle et R₁ est le groupe méthyle; et sels d'addition pharmaceutiquement acceptables de celui-ci.


2. Composé selon la revendication 1, dans lequel R représente le groupe méthoxyméthyle et les autres substituants sont tels que définis dans la revendication 1.

3. Composé selon la revendication 1, dans lequel R représente le groupe méthoxyméthyle, R₁ et R₄ représentent chacun le groupe méthyle et Y représente un groupe de formule



dans laquelle R₂ est un atome d'hydrogène et R₃ est tel que défini dans la revendication 1.

4. Composé selon la revendication 1, dans lequel R est le groupe méthoxyméthyle, R₁ et R₄ sont chacun le groupe méthyle et Y est un fragment aminé qui dérive d'un aminoacide naturel tel que la glycine, l'ornithine, la sérine, l'acide aspartique, la tyrosine, la leucine, la phénylalanine, la méthionine, la proline, la thréonine, la lysine, ou d'un dipeptide synthétique tel que la glycyllysine, la sérylproline, le glycylprolinamide, le tyrosylprolinamide, le thréonylprolinamide, le leucylprolinamide.
5. Composé selon la revendication 1, dans lequel R est le groupe méthoxyméthyle, R₁ et R₄ sont chacun le groupe méthyle, Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est une chaîne alkyle linéaire ayant de préférence de 3 à 12 atomes de carbone, encore mieux de 3 à 7 atomes de carbone, substituée par un groupe choisi parmi COOH, SO₃H et PO₃H₂.
6. Composé selon la revendication 1, dans lequel R est le groupe méthoxyméthyle, R₁ et R₄ sont chacun le groupe méthyle, Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène, et R₃ est CH₂CH₂CH₂CH₂CH₂-COOH.
7. Composé selon la revendication 1, dans lequel R représente un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ représente un atome d'hydrogène ou le groupe méthyle, R₄ représente un atome d'hydrogène, le groupe méthyle ou hydroxyméthyle, et Y représente un groupe de formule



dans laquelle R₂ est un atome d'hydrogène et R₃ est tel que défini dans la revendication 1.

8. Composé selon la revendication 7, dans lequel Y est un fragment aminé qui dérive d'un aminoacide naturel tel que la glycine, l'ornithine, la sérine, l'acide aspartique, la tyrosine, la leucine, la phénylalanine, la méthionine, la proline, la thréonine, la lysine, ou d'un dipeptide synthétique tel que la glycyllysine, la sérylproline, le glycylprolinamide, le tyrosylprolinamide, le thréonylprolinamide, le leucylprolinamide.
9. Composé selon la revendication 1, dans lequel R est un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ est un atome d'hydrogène ou le groupe méthyle, R₄ est un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle, et Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est une chaîne alkyle linéaire ayant de préférence 3 à 12 atomes de carbone, encore mieux de 3 à 7 atomes de carbone, substituée par un groupe choisi parmi COOH, SO₃H et PO₃H₂.
10. Composé selon la revendication 1, dans lequel R est un atome d'hydrogène ou le groupe hydroxyméthyle ou méthoxyméthyle, R₁ est un atome d'hydrogène ou le groupe méthyle, R₄ est un atome d'hydrogène ou le groupe méthyle ou hydroxyméthyle, et Y est un groupe NR₂R₃ dans lequel R₂ est un atome d'hydrogène et R₃ est CH₂CH₂CH₂CH₂CH₂-COOH.

C₄)carbonyle, mercapto, alkyl(C₁-C₄)thio dans lequel le fragment alkyle peut être éventuellement substitué par un groupe carboxy, phényle qui peut être éventuellement substitué par 1 à 3 substituants choisis parmi des groupes carboxy, sulfo, hydroxy, halogéno

5 et mercapto, carbamoyle, alkyl(C₁-C₆)carbamoyle dans lequel le fragment alkyle peut être éventuellement substitué par 1 ou 2 substituants choisis parmi des groupes carboxy, amino, alkyl(C₁-C₄)amino et dialkyl-(C₁-C₄)amino, dialkyl(C₁-C₄)-carbamoyle dans lequel les fragments alkyle conjointement avec l'atome d'azote adjacent peuvent également représenter un cycle hétérocyclique saturé à 5-7 chaînons qui peut éventuellement être substitué par un groupe carboxy ou carbamoyle sur l'un des atomes de carbone formant le cycle, et qui peut éventuellement contenir un autre hétéroatome
10 choisi parmi O, S et N, benzoylamino dans lequel le fragment phényle peut être substitué par 1 à 3 groupes hydroxy, un cycle hétérocyclique azoté contenant 5-6 chaînons, qui peut être insaturé, partiellement saturé ou totalement saturé et peut contenir 1 à 3 autres hétéroatomes choisis parmi N, S et O, l'un des atomes de carbone du cycle pouvant éventuellement porter un groupe carboxy, sulfo, carboxy-alkyle(C₁-C₄) ou sulfo-alkyle(C₁-C₄) et l'atome d'azote du cycle pouvant éventuellement être substitué par un groupe alkyle en C₁-C₄, carboxy-alkyle(C₁-C₄), sulfo-alkyle(C₁-C₄) ou benzyle;

20 un groupe alcényle en C₃-C₆, éventuellement substitué par le groupe carboxy ou sulfo; le groupe 1-désoxy-1-glucityle; le groupe 2-désoxy-2-glucosyle; un cycle hétérocyclique azoté à 5-7 chaînons, totalement saturé, dans lequel l'atome d'azote peut être éventuellement substitué par un groupe alkyle en C₁-C₄ ou benzyle, et 1 ou 2 atomes de carbone du squelette cyclique peuvent porter un substituant choisi
25 parmi des groupes alkyle en C₁-C₄, carboxy et sulfo; ou

R₂ et R₃ forment ensemble, avec l'atome d'azote adjacent, un cycle hétérocyclique à 5-7 chaînons totalement saturé, qui peut éventuellement contenir un autre hétéroatome
30 choisi parmi O, S et N, et peut éventuellement porter 1 ou 2 substituants sur les atomes de carbone formant le cycle, choisis parmi des groupes alkyle en C₁-C₄, benzyle, carboxy, sulfo, carboxy-alkyle(C₁-C₄) et sulfo-alkyle(C₁-C₄);

R₄ représente un atome d'hydrogène,
le groupe méthyle ou
35 le groupe hydroxyméthyle,

sous réserve que lorsque R₄ est un atome d'hydrogène ou le groupe hydroxyméthyle, R est alors simultanément le groupe méthoxyméthyle et R₁ est le groupe méthyle; et de ses sels d'addition pharmaceutiquement acceptables, qui comprend la mise en réaction d'un composé antibiotique GE 2270 de formule II:

40

45

50

55



30

- 35

40

- 45

- 50

- 55